



THE CROATIAN CHEMICAL SOCIETY
PROUDLY PRESENTS

THE COMPILATION OF THIS YEAR'S NEW

SOLUTIONS IN CHEMISTRY

8-11 November 2022 • Sveti Martin na Muri, Croatia

INTERNATIONAL CONFERENCE RECOGNISED BY THE EUROPEAN CHEMICAL SOCIETY (EuChemS)



BOOK OF ABSTRACTS



SOLUTIONS IN CHEMISTRY 2022

8–11 November 2022

Hotel Terme Sveti Martin☆☆☆☆

Sveti Martin na Muri, Croatia

BOOK OF ABSTRACTS

IMPRESSUM

ORGANIZER

Croatian Chemical Society

PUBLISHED BY

Croatian Chemical Society
Horvatovac 102a, 10000 Zagreb, Croatia

EDITORS

Petar Kassal, Ernest Meštrović, Danijel Namjesnik, Rosana Ribić,
Marina Šekutor, Vladislav Tomišić, Andrea Usenik

CONFERENCE VENUE

Hotel Terme Sveti Martin^{☆☆☆☆}
Izvorska 3, 40313 Sveti Martin na Muri, Croatia
www.termesvetimartin.com

solutionsinchemistry.hkd.hr
sic@hkd.hr

ISBN 978-953-8334-05-4

Zagreb, 2022

UNDER THE AUSPICES OF

Zoran Milanović, President of the Republic of Croatia

Ministry of Economy and Sustainable Development of the Republic of Croatia
Ministry of Science and Education of the Republic of Croatia

Croatian Academy of Sciences and Arts
Croatian Academy of Engineering

Ruđer Bošković Institute

University of Zagreb
University North

Croatian Chamber of Economy
Croatian Employers' Association
Croatian Engineering Association
Croatian Society for Natural Sciences

Faculty of Science (UniZg)
Faculty of Chemical Engineering and Technology (UniZg)

Teacher Education and Training Agency, Croatia

City of Čakovec
City of Mursko Središće

SPONSORS

(IN ALPHABETICAL ORDER)

GOLDEN SPONSORS



SILVER SPONSORS



BRONZE SPONSORS



EXHIBITORS



SUPPORTERS AND DONORS



MEDIA SPONSOR



ORGANISATION

SCIENTIFIC AND ORGANISING COMMITTEE

Ernest Meštrović, *Chair*

Vladislav Tomišić, *Chair*

Andrea Usenik, *secretary*

Klemen Bohinc

Nikola Bregović

Predrag Čudić

Petar Kassal

Danijel Namjesnik

Predrag Novak

Josip Požar

Rosana Ribić

Giovanna Speranza

Marina Šekutor

Ines Topalović Piteša

EDUCATION SUBCOMMITTEE

Olgica Martinis, *Chair*

Jurica Bauer

Dražen Crnčec

Vladimir Stilinović

Welcome from the Chairmen of the Organising Committee

On behalf of the Organising Committee, it is our great pleasure to welcome you to the Solutions in Chemistry 2022 conference taking place in Sveti Martin na Muri from 8 to 11 November 2022. The focus of the Conference organised by the Croatian Chemical Society can be deduced from the two meanings of the word “Solutions” in its title. The first one refers to the solution as a phase and comprises the experimental and computational studies of the processes occurring in liquid media. The second one is concerned with general solutions to chemical problems which are met in academia and various branches of industry. Thus, the Conference aims to gather the scientists dealing with different fields of chemistry and to deepen and strengthen the collaboration of the experts from universities, research institutes, schools, and industrial companies.

A number of plenary, invited, and section lectures will be delivered by top-tier scientists, and various sponsor presentations and workshops will be held as well. These will provide a new insight into the current problems in chemical science and industry, and an opportunity to learn and obtain new skills from prominent chemists and entrepreneurs. In addition, the Conference participants will have a chance to find out more about the newest instruments and consumables available from our numerous sponsors, exhibitors, and donors to whom we are grateful for their support. The poster session will give everyone the opportunity to present their work: from undergraduate students to experienced scientists in diverse fields of chemistry and chemical engineering. The education section will be held in parallel, focusing on finding new solutions in teaching new generations of chemists.


A significant emphasis will also be put on an irreplaceable part of every great conference: social activities. Thus, we plan to organize various social events and encourage the participants to interact in a non-formal manner as well as to take the opportunity to discover the beauty of the picturesque Međimurje County and taste the local delicacies.

LET'S FIND THE NEW SOLUTIONS IN CHEMISTRY!

Ernest Meštrović & Vladislav Tomišić

PLENARY LECTURERS

(IN CHRONOLOGICAL ORDER)




Professor **Werner M. Nau** obtained his MSc in 1991 at St. Francis Xavier University, Canada, and his PhD in 1994, at the University of Würzburg, Germany. He was a postdoctoral researcher at the University of Ottawa, Canada, and became an Assistant Professor at the University of Basel, Switzerland. Since 2002, he is a Professor of Chemistry at Jacobs University Bremen (Germany), and, since 2011, Dean of the faculty. Prof. Nau has held visiting professorships at the University of Wisconsin-Madison (USA), the University of Cambridge (UK), the University of Huelva (Spain), and the Institute of Chemical Research of Catalonia (Spain). His research combines principles of organic photochemistry with spectroscopy and supramolecular chemistry and is directed towards applications in bioanalytical chemistry, peptide folding, drug delivery, drug discovery, catalysis, chiral recognition, fluorescent dye design, chemosensing, as well as enzyme and membrane assays. His research is published in ca. 300 research publications, 6 book chapters, and 4 patents. Prof. Nau received several scientific awards; the Grammaticakis–Neumann Prize of the European Photochemistry Association (2000), the ADUC-Jahrespreis für Habilitanden (2000), the Werner Prize 2002 of the Swiss Chemical Society, and the EPA-PPS Prize 2010. He has served, among others, as President (2012–2014) of the European Photochemistry Association (EPA) and Vice Chairman (2010–2013) of the Photochemistry Section of the German Chemical Society.




Dr **Aleksandar Danilovski** obtained his MSc in chemistry in 1997 at the Faculty of Science, University of Zagreb, and his PhD in 2001 at the same faculty. He started his career in the industry at the PLIVA Research Institute, where he was a member of the Management Board for 20 years. In 2009 he held the position of Senior Vice President of the Management Board XELLIA Pharmaceuticals (Denmark), and the position of Chief Scientific Officer (CSO), which he still holds today. In 2011, he founded the research company XELLIA Ltd. (Croatia) where he is the Managing Director. Dr Danilovski is one of the co-founders of a biotechnology start-up company (Pharmaero, Denmark). He is a member of the Board of Directors in Pharmaero (Denmark), of the Scientific Selection Board in Novo Holdings, REPAIR Impact Fund (Denmark), of the Board of Directors in Active Biotech (Sweden), and of the Scientific Advisory Board in Bugworks (USA/India). In 2017, he was elected as a member of the Committee for Chemistry of the Croatian Academy of Sciences and Arts, and since 2018 he is a member of the Croatian Academy of Engineering. Dr Danilovski led the research and development, registration and launch on the global market (US and EU) of more than 100 different active substances and over 200 different finished pharmaceutical products of which several innovative and patent-protected medicines with added value.

PLENARY LECTURERS

(IN CHRONOLOGICAL ORDER)




Professor **Eric Bakker** was educated at the Swiss Federal Institute of Technology (ETH) in Zurich, Switzerland and earned his PhD in 1993. He spent many years in the United States and Australia before moving to the University of Geneva in Switzerland where he is a professor of analytical chemistry. His research interests are in the area of chemical sensors for environmental (aquatic) and biomedical applications. The main emphasis is the development and understanding of electrochemical and optical sensing principles, often using membrane transport, complexation, and extraction phenomena. New principles and materials are developed, prototype instrumentation is engineered, and their use is tested in the final sample environment of interest, making this research quite broad in scope. His research group has published nearly 400 highly cited research papers. Prof. Bakker also serves as the Executive Editor of the journal ACS Sensors.




Professor **Csaba Szántay, Jr.** received his MSc in chemical engineering at the Technical University of Budapest in 1982, where he obtained his PhD in 1986. He was a postdoctoral fellow at Leeds University (UK) in 1988, and after he continued his work as an NMR spectroscopist in the Spectroscopic Research Department of Gedeon Richter Plc., Hungary's largest pharmaceutical company. Since 1994 he has been the Head of this department. He and his team are committed to maintaining strong ties with academia in terms of education as well as research collaborations. Prof. Szántay habilitated at the TUB in 2002, and in 2003 elected to be a Private Professor at the same university. His main fields of research are the physical and mathematical theory of NMR spectroscopy and the application of high-resolution NMR in the structure determination of organic molecules. He obtained from the Hungarian Academy of Sciences a Candidature in 1991 and a Doctor of Sciences title in 2000. One major project of his that transpired from blending philosophy, psychology, physics, mathematics, and applied NMR and MS, was published worldwide as a book (*Anthropic Awareness: The Human Aspects of Scientific Thinking in NMR Spectroscopy and Mass Spectrometry*. New York: Elsevier, 2015.). He has published around 190 original scientific papers, including three books. He also holds numerous high-ranking positions in various scientific committees and institutions, as well as in several foundations aimed at supporting the teaching of natural sciences in elementary and secondary schools.

PLENARY LECTURERS

(IN CHRONOLOGICAL ORDER)



Professor **Jurij Lah** obtained his PhD in Chemistry in 2000 at the Faculty of Chemistry and Chemical Technology, University of Ljubljana (UL FCCT), Slovenia. He was a postdoctoral researcher at the Vrije Universiteit Brussel (VUB), Belgium. Since 2013, he is a full professor in charge of courses Physical Chemistry, Biophysical Chemistry, Chemical Thermodynamics and Biomolecular Thermodynamics at the Faculty of Chemistry and Chemical Technology, University of Ljubljana, Slovenia. Prof. Lah is a distinguished lecturer, awarded four times (in 2004, 2011, 2018, and 2020) for the best lecturer at the UL FCCT. He is very active in the popularisation of science through interviews, lectures, and cover pages. He is currently principal investigator in several basic research projects and projects with the pharmaceutical industry. His main research areas include different topics related to thermodynamic stability, interactions, and structural features of biologically important molecules in relation to their function, and his recent research is focused on describing the behaviour of non-canonical nucleic acids and intrinsically disordered proteins. Prof. Lah publishes research results in reputable journals (*PNAS*, *Angew. Chem.*, *JACS*, *Nucleic Acids Res.*, etc.). He was awarded in 2014 for outstanding research achievements in chemistry and related sciences, and in 2019 he received Zois Award for Outstanding Achievements in Physical Biochemistry.



Professor **Angela Danil de Namor** is the first female professor of chemistry at the University of Surrey, UK. She graduated in biochemistry at the National University of the South, Bahia Blanca, Argentina and in 1973 obtained a PhD in chemistry at the University of Surrey, UK. After postdoctoral research sponsored by Unilever Research Laboratories, she joined the Department of Chemistry in 1977 where she is now an emeritus professor of physical chemistry. Her research interest is in the area of solution thermodynamics involving mostly supramolecular systems; solution thermodynamics of electrolytes, cyclodextrins and calix-based receptors, their interaction with ionic and neutral species and their broad applications. She published more than 170 papers in prestigious chemical journals and supervised over 50 PhD students. She has coordinated several EU contracts involving Partners from Europe, Latin America, Africa, and Middle East Countries. Prof. Danil de Namor has received numerous awards and recognitions; the Doctor Honoris Causa from three Universities in South America for contributions given to the development of chemistry in Latin America, fellow of the Royal Society of Chemistry and fellow of the International Union of Pure & Applied Chemistry (IUPAC), emeritus fellowship from the Leverhulme Trust (2014). She was the first foreign woman elected member of the National Academy of Sciences of Argentina and in 2021 Prof. Danil de Namor was awarded the Doctor of Science degree by the University of Surrey for proven evidence of the international impact of her research to advances in the discipline, following the thesis entitled “Supramolecular Chemistry: Personal Contributions to the Discipline”.

PLENARY LECTURERS

(IN CHRONOLOGICAL ORDER)



Dr **Nicoletta Ravasio** graduated in chemistry at the University of Milano (Italy) in 1982, and in 1987 obtained PhD in chemical sciences at the University of Bari, Italy. She is currently the research director at the “Giulio Natta” Institute of Chemical Sciences and Technologies (SCITEC) of the National Research Council (CNR), Italy. Her main scientific interests are sustainable organic synthesis and renewable raw materials. In the field of vegetable oils, she developed a selective hydrogenation process, a process for the esterification and transesterification and a process for the synthesis of bio-lubricants. Recently she has been devoted to the valorization of agri-food waste with different added value, from building materials to cosmetics, performing even better than the fossil-based ones. Dr Ravasio is principal investigator in several national and international research projects and contracts, and the author of highly cited 164 papers and 6 patents. As a recognized expert she evaluated EU Horizon 2020-Waste 7 and Horizon2020-Bio Based Industries projects. Since 2015 she is chair of Oleochemistry, Molecule and Polymer Science division of European Federation on Lipid Science, and since 2018 a member of Technical-Scientific Committee Italian Cluster of Green Chemistry (SPRING), and a member of Team of Experts to support the High Level Roundtable on the implementation of the Chemicals Strategy for Sustainability (EuChemS). In 2021 she was awarded with the Piero Pino medal of the Italian Chemical Society "For studies of scientific, innovative and applicative relevance in the field of heterogeneous catalysis with particular attention to the substitution of noble metals in fine chemicals synthesis and to selective transformation of renewable raw materials".

INVITED LECTURES

(IN CHRONOLOGICAL ORDER)

PREDRAG ČUDIĆ
Charles E. Schmidt College of Science,
Florida Atlantic University, Boca Raton, USA

POTENTIAL SOLUTION FOR PEPTIDE-BASED DRUG DELIVERY TO THE BRAIN: DESIGN AND ACTIVITY OF MULTIFUNCTIONAL CYCLIC PEPTIDE-BASED ANALGESICS



INVITED LECTURE
DAY 1
TUESDAY, 8 November



TOMISLAV BILJAN
PLIVA Croatia Ltd., Zagreb, Croatia

NMR SPECTROSCOPY AND GENERIC PHARMA R&D: PAST, PRESENT AND THE FUTURE



INVITED LECTURE
DAY 1
TUESDAY, 8 November



LUNA MASLOV BANDIĆ
Faculty of Agriculture, University of Zagreb, Croatia

MOLECULES BEHIND WINE TASTING



INVITED LECTURE
DAY 1
TUESDAY, 8 November



IVA REZIĆ
Faculty of Textile Technology, University of Zagreb, Croatia

SOLVING COMPLEX PROBLEMS IN SOLUTION CHEMISTRY BY DESIGN OF EXPERIMENT



INVITED LECTURE
DAY 2
WEDNESDAY, 9 November



EWALD EDINK
Holland University of Applied Sciences, Amsterdam, The Netherlands

SUPPORTING STUDENTS IN SOLVING QUANTITATIVE CHEMICAL PROBLEMS USING AN ADAPTIVE DIGITAL PLATFORM



INVITED LECTURE
DAY 2
WEDNESDAY, 9 November



KREŠIMIR MOLČANOV
Ruder Bosković Institute, Zagreb, Croatia

CHEMISTRY IN THREE DIMENSIONS: CHIRALITY, SYMMETRY, ISOMERISM



INVITED LECTURE
DAY 2
WEDNESDAY, 9 November



NIKOLINA LETIĆ
Lumménice Ltd., Zagreb, Croatia

FUNDING OPPORTUNITIES IN INNOVATIVE PROJECTS FOR THE 2021–2027 PERIOD



INVITED LECTURE
DAY 2
WEDNESDAY, 9 November



VLADIMIR STILINOVIĆ
Faculty of Science, University of Zagreb, Croatia

DISCOVERIES, CONTROVERSIES AND BEHOLDINGS – HOW WATER STOPPED BEING AN ELEMENT



INVITED LECTURE
DAY 2
WEDNESDAY, 9 November



ŽIGA JAKOPIN
Faculty of Pharmacy, University of Ljubljana, Slovenia

STRUCTURAL REQUIREMENTS FOR IN VITRO AND IN VIVO ACTIVITY OF NOD2 AGONISTS



INVITED LECTURE
DAY 3
THURSDAY, 10 November



GIUSEPPE CAPPELLETTI
Department of Chemistry, University of Milan, Italy

TAILORING THE LIQUID/LIQUID INTERFACE BY TRADITIONAL AND PICKERING GREEN EMULSIONS



INVITED LECTURE
DAY 3
THURSDAY, 10 November



N A G R A D A
Leopold Ružička

Winner of the
Leopold Ružička Award
given by the
Croatian Chemical Society

IGOR ŽIVKOVIĆ
Faculty of Science, University of Zagreb, Croatia

ALL BUT ONE: HOW NEGATIVE CATALYSIS SHAPED EVOLUTION OF THE ISOLEUCYL-tRNA SYNTHETASE



LEOPOLD RUŽIČKA AWARD WINNER
INVITED LECTURE
DAY 4
FRIDAY, 11 November



CONFERENCE PROGRAMME

DAY 1 • TUESDAY, 8 November

9:00 – 14:00	REGISTRATION EARLY CHECK IN from 12 h for guests of Hotel Terme Sveti Martin	 PRE-CONFERENCE WORKSHOP (Mura III)	
		9:30 – 10:30	Solubility Crystallization with Small Scale Parallel Crystallizers
		10:30 – 10:45	BREAK
		10:45 – 11:45	API Improvements through Co-Crystallization
		11:45 – 12:00	BREAK
		12:00 – 12:30	Nucleation and Induction Time Measurements
14:00 – 14:40	OPENING CEREMONY (Mura Halls)		
	SECTION I (Mura Halls) (Chair: Vladislav Tomišić)		
14:40 – 15:25	PLENARY	Werner M. Nau: SOLUTIONS IN SUPRAMOLECULAR CHEMISTRY IN WATER	
15:25 – 15:55	INVITED	Predrag Čudić: POTENTIAL SOLUTION FOR PEPTIDE-BASED DRUG DELIVERY TO THE BRAIN: DESIGN AND ACTIVITY OF MULTIFUNCTIONAL CYCLIC PEPTIDE-BASED ANALGESICS	
15:55 – 16:15	SECTION	Gordan Horvat: Cyclopeptides as Versatile Anion Sensors in Solution	
16:15 – 16:35		Gracjan Kurpik: Metallosupramolecular Assemblies – From Simple Complexes to Functional Nanostructures	
16:35 – 17:15	☕ COFFEE BREAK (in front of Mura Halls) ☕		
	SECTION II (Mura Halls) (Chair: Ernest Meštrović)		
17:15 – 18:00	PLENARY	Aleksandar Danilovski: CORRELATION DOES NOT NECESSARILY IMPLY CAUSATION	
18:00 – 18:30	INVITED	Tomislav Biljan: NMR SPECTROSCOPY AND GENERIC PHARMA R&D: PAST, PRESENT AND THE FUTURE	
18:30 – 19:00	INVITED	Luna Maslov Bandić: MOLECULES BEHIND WINE TASTING	
19:00 – 20:15	🍷 DINNER BREAK 🍷		
20:15 – 21:15	WINE TASTING by Štampar Winery (Restaurant Falat)		
21:15 – 22:45	WELCOME PARTY (Restaurant Falat)		

DAY 2 • WEDNESDAY, 9 November

8:30 – 9:45	REGISTRATION FOR TEACHERS (1 DAY)	
	SECTION III – PART ONE (Mura Halls) (Chair: Petar Kassal)	
9:00 – 9:45	PLENARY	Eric Bakker: SENSING SOLUTIONS
	EDUCATION SECTION – PART ONE (Chair: Vladimir Stilinović)	
9:45 – 10:30	PLENARY	Csaba Szántay, Jr.: THE HUMAN FACTOR IN SCIENTIFIC THINKING: THE ILLUSIONS THAT WE LIVE BY
10:30 – 11:00	 COFFEE BREAK (Ion Bar) 	

MORNING SECTIONS:

	SECTION III – PART TWO (Mura I + II) (Chair: Petar Kassal)	
11:00 – 11:20	SECTION	Ivo Piantanida: Triarylboranes: Novel Type of Dye for Biomacromolecules
11:20 – 11:40		Miroslava Čonková: From Single Molecule to Responsive Aggregate: Multiresponsive Photoswitch in Nonpolar Solvents
11:40 – 12:00		Nikola Sakač: Solutions for Pollutants Sensing – Surfactants, Estrogen and Glyphosate Sensors
12:00 – 12:30	SPONSORS	Selvita
12:30 – 12:40		CAS
12:40 – 12:50		Labtim
12:50 – 13:00		Shimadzu

	EDUCATION SECTION – PART TWO (Mura III) (Chairs: Jurica Bauer, Vladimir Stilinović)	
11:00 – 11:30	INVITED	Ewald Edink: SUPPORTING STUDENTS IN SOLVING QUANTITATIVE CHEMICAL PROBLEMS USING AN ADAPTIVE DIGITAL PLATFORM
11:30 – 12:00	INVITED	Krešimir Molčanov: CHEMISTRY IN THREE DIMENSIONS: CHIRALITY, SYMMETRY, ISOMERISM
12:00 – 12:20	SECTION	Mirta Malčić, Suzana Lovrić: From Old to New Towards Sustainable Development
12:20 – 12:40		Daria Stejskal: Innovative Approaches in Teaching Chemistry
12:40 – 13:00		Nikola Bedeković: Carbonated Water: Acid, Buffer or Both?
13:00 – 14:30	 LUNCH BREAK 	

AFTERNOON SECTIONS:

14:30 – 15:30	SIC × AlphaChrom PUB QUIZ – details at the registration desk!	
15:30 – 15:50	SHORT BREAK	
	SECTION IV (Mura I + II) (Chair: Ines Topalović Piteša)	
15:50 – 16:20	INVITED	Iva Rezić: SOLVING COMPLEX PROBLEMS IN SOLUTION CHEMISTRY BY DESIGN OF EXPERIMENT
16:20 – 16:50	INVITED	Nikolina Letić: FUNDING OPPORTUNITIES IN INNOVATIVE PROJECTS FOR THE 2021–2027 PERIOD
16:50 – 17:30	POSTER SETUP	
	EDUCATION SECTION – PART THREE (Mura III) (Chair: Vladimir Stilinović)	
14:30 – 14:50	SECTION	Marijana Bastić: How to Facilitate the Evaluation of Student Achievements by Using Written Papers in Chemistry?
14:50 – 15:10		Monika Pavić, Maja Jurić-Babaj: Challenges in Chemistry Classes Using ICT Technology
15:10 – 15:30		Jurica Bauer: Fostering Transdisciplinary Learning in Science and Technology
15:30 – 15:50		Olgica Martinis: The Current Challenges of Improving Chemistry Learning
15:50 – 17:00	ROUND TABLE	What Our Education Gave Us? (“Što je nama naša nastava dala”) – Mementos, comments, and critiques by former Croatian students (Filip Topić, Ilija Čorić, Jana Volarić, Vedran Vuković, Petra Vizjak, Anđela Šarić)
17:00 – 17:30	EVALUATION	
17:30 – 20:00	POSTER SESSION (Mura I + II) BEER BREAK by local craft brewery “Lepi Dečki”	
20:00 – 21:00	 DINNER BREAK 	
	SECTION H₂O (Pub Potkova) (Chair: Nikola Bregović)	
21:00 – 21:30	INVITED	Vladimir Stilinović: DISCOVERIES, CONTROVERSIES AND BEHEADINGS – HOW WATER STOPPED BEING AN ELEMENT
21:30 – ... ☺	#MoreThanWater CONFERENCE PARTY with DJ Bregi	

DAY 3 • THURSDAY, 10 NOVEMBER

SECTION V – PART ONE (Mura I + II) (Chair: Rosana Ribić)		
9:00 – 9:45	PLENARY	Jurij Lah: ADVENTURES IN PHASE SPACE OF DNA BINDING AND STABILITY
9:45 – 10:15	INVITED	Žiga Jakopin: STRUCTURAL REQUIREMENTS FOR IN VITRO AND IN VIVO ACTIVITY OF NOD2 AGONISTS
10:15 – 10:25	SPONSORS	IMROH (IMI)
10:25 – 10:35		Jasika
10:35 – 11:10	 COFFEE BREAK (in front of Mura Halls) 	
SECTION V – PART TWO (Mura I + II) (Chair: Rosana Ribić)		
11:10 – 11:30	SECTION	Maša Safundžić Kučuk: JGL's Experience in Demonstrating Extended Pharmaceutical Equivalence in Drug Product Solutions
11:30 – 11:50		Želimir Kurtanjek: Causal Analysis of Molecular Descriptors of Eutectic Solutions of Choline Chloride Systems Applicable in Pharma Technologies
11:50 – 12:10		Riya Sailani: Spectrophotometric Kinetic and Thermodynamic Investigation for the Oxidation of Anilines by Hexachloroiridate(IV) in Solution Media: A Combined Experimental and Computational Study
12:10 – 13:45	 LUNCH BREAK 	
SECTION VI (Mura I + II) (Chair: Josip Požar)		
13:45 – 14:30	PLENARY	Angela Danil de Namor: SUPRAMOLECULAR CHEMISTRY: SOLUTION THERMODYNAMICS AND APPLICATIONS
14:30 – 15:00	INVITED	Giuseppe Cappelletti: TAILORING THE LIQUID/LIQUID INTERFACE BY TRADITIONAL AND PICKERING GREEN EMULSIONS
15:00 – 15:20	SECTION	Aleks Logožar: ML ₂ Metal Complexes with bpa and imda Ligands – In Solution and Solid-State
15:20 – 15:40		Marin Liović: Halogenide Anions as Hydrogen and Halogen Bond Acceptors in Simple Cocrystals with a Neutral Halogen Bond Donor
CONFERENCE EXCURSION		
16:15 – 17:00	ORGANISED BUS TRANSIT: Terme Sveti Martin → Čakovec + a surprise visit ... The buses (GROUP 1 and GROUP 2) leave at 16:15 from the main entrance to Hotel Terme Sveti Martin. PLEASE BE THERE AT 16:00. Thank you!	
17:00 – 18:15	GROUP 1 Treasury of Međimurje	GROUP 2 Local food tasting by Slađana Herman
18:15 – 18:30	GROUP SWITCH	
18:30 – 19:45	GROUP 1 Local food tasting by Slađana Herman	GROUP 2 Treasury of Međimurje
19:45 – 20:15	ORGANISED BUS TRANSIT: Čakovec → Terme Sveti Martin	
21:00 – ... 	CONFERENCE DINNER (Mura Halls) Classical serving, local food paired with local wines, live music	

DAY 4 • FRIDAY, 11 NOVEMBER

9:00 – 9:30	WAKE UP LECTURE	Surprise, surprise ... A positive one!	
	SECTION VII (Mura I + II) (Chair: Giovanna Speranza)		
9:30 – 10:15	PLENARY	Nicoletta Ravasio: CHEMICAL SOLUTIONS FOR AGRI–FOOD WASTE UPCYCLING	
10:15 – 10:35	SECTION	Davor Kovačević: Applications of Polyelectrolyte Multilayers for the Reduction of Bacterial Adhesion	
10:35 – 10:55		Ina Erceg: Comparison of Albumin and Chitosan Effects on Calcium Phosphate Formation on Titanate Nanomaterials	
10:55 – 11:15		Gloria Zlatić: Protective Ability of <i>Artemisia Annu</i> L. Against Microbiologically Influenced Corrosion of Aluminium Alloy 5083 Caused by <i>Pseudomonas Aeruginosa</i> in Artificial Seawater	
	SECTION VIII (Mura I + II) (Chair: Marina Šekutor)		
11:15 – 11:45	INVITED Winner of the Leopold Ružička Award given by CCS	Igor Živković: ALL BUT ONE: HOW NEGATIVE CATALYSIS SHAPED EVOLUTION OF THE ISOLEUCYL–tRNA SYNTHETASE’S EDITING DOMAIN	
11:45 – 12:00	BEST POSTER PRESENTATION Winner of the Tomislav Cvitaš Award given by CCS	Presenting author to be announced after the Poster session ...	
12:00 – 12:30	CLOSING CEREMONY SEE YOU IN 2024!		

CONTENTS

PLENARY LECTURES

PL1	Werner M. Nau Solutions in Supramolecular Chemistry in Water	24
PL2	Eric Bakker, Tara Forrest, Elena Zdrachek, Polyxeni Damala, Pitchnaree Kraikaew, Robin Nussbaum Sensing Solutions With Potentiometric Probes	25
PL3	Csaba Szántay The Human Factor in Scientific Thinking: The Illusions That We Live By	27
PL4	Jurij Lah, San Hadži Adventures in Phase Space of Dna Binding and Stability	28
PL5	Angela F. Danil de Namor Supramolecular Chemistry: Solution Thermodynamics and Its Applications	31
PL6	Nicoletta Ravasio Chemical Solutions for Agri-Food Waste Upcycling	32

INVITED LECTURES

IL1	Predrag Cudic, Shahayra Majumder, Shahayra Majumder, Heather Stacy, Thomas Cirino, Brandon Williams, Jay P. McLaughlin Potential Solution for Peptide-Based Drug Delivery to the Brain: Design and Activity of Multifunctional Cyclic Peptide-Based Analgesics	36
IL2	Tomislav Biljan NMR Spectroscopy and Generic Pharma R&D: Past, Present and the Future	37
IL3	Luna Maslov Bandić Molecules Behind Wine Tasting	39
IL4	Iva Rezić Solving Complex Problems in Solution Chemistry by Design of Experiment	40
IL5	Ewald Edink, Jeroen Bottema, Artur Lorincz, Kinga Lorincz The Path Towards Enlightenment? Supporting Students in Solving Complex Chemical Calculations Using an Adaptive Digital Platform	42
IL6	Krešimir Molčanov Chemistry in Three Dimensions: Chirality, Symmetry, Isomerism	43
IL7	Nikolina Letić Funding Opportunities in Innovative Projects for the 2021–2027 Period	45
IL8	Vladimir Stilinović Discoveries, Controversies and Beheadings – How Water Stopped Being an Element	46
IL9	Žiga Jakopin Structural Requirements for <i>In Vitro</i> and <i>In Vivo</i> Activity of NOD2 Agonists	48
IL10	Giuseppe Cappelletti Tailoring the Liquid/Liquid Interface of Traditional and Pickering Green Emulsions	49
IL11	Igor Živković, Aleksandra Maršavelski, Ita Gruić Sovulj All But One: How Negative Catalysis Shaped Evolution of the Isoleucyl-tRNA Synthetase's Editing Domain	51

SECTION LECTURES

SL1	Gordan Horvat Cyclopeptides as Versatile Anion Sensors in Solution	53
SL2	Gracjan Kurpiak Metallosupramolecular Assemblies – From Simple Complexes to Functional Nanostructures.....	54
SL3	Miroslava Čonková From Single Molecule to Responsive Aggregate: Multiresponsive Photoswitch in Nonpolar Solvents.....	56
SL4	Nikola Sakač Solutions for Pollutants Sensing – Surfactants, Estrogen and Glyphosate Sensors	57
SL5	Ivo Piantanida Triarylboranes: Novel Type of Dye for Biomacromolecules	59
SL6	Maša Safundžić Kučuk JGL's Experience in Demonstrating Extended Pharmaceutical Equivalence in Drug Product Solutions.....	60
SL7	Želimir Kurtanjek Causal Analysis of Molecular Descriptors of Eutectic Solutions of Choline Chloride Systems Applicable in Pharma Technologies	62
SL8	Riya Sailani Spectrophotometric Kinetic and Thermodynamic Investigation for the Oxidation of Anilines by Hexachloroiridate(IV) in Solution Media: A Combined Experimental and Computational Study	63
SL9	Aleks Logožar ML ₂ Metal Complexes with bpa and imda Ligands – In Solution and Solid-State	65
SL10	Marin Liović Halogenide Anions as Hydrogen and Halogen Bond Acceptors in Simple Cocrystals with a Neutral Halogen Bond Donor.....	66
SL11	Davor Kovačević Applications of Polyelectrolyte Multilayers for the Reduction of Bacterial Adhesion	68
SL12	Ina Erceg Comparison of Albumin and Chitosan Effects on Calcium Phosphate Formation on Titanate Nanomaterials.....	69
SL13	Gloria Zlatić Protective Ability of <i>Artemisia Annu</i> L. Against Microbiologically Influenced Corrosion of Aluminium Alloy 5083 Caused by <i>Pseudomonas Aeruginosa</i> in Artificial Seawater	70

EDUCATION SECTION LECTURES

ESL1	Daria Stejskal Innovative Approaches in Teaching Chemistry	71
ESL2	Mirta Malčić, Suzana Lovrić From Old to New Towards Sustainable Development.....	72
ESL3	Nikola Bedeković Carbonated Water: Acid, Buffer or Both?	73
ESL4	Monika Pavić, Maja Jurić-Babaja Challenges in Chemistry Classes Using ICT Technology.....	74

ESL5	Marijana Bastić How to Facilitate the Evaluation of Student Achievements by Using Written Papers in Chemistry?	75
ESL6	Jurica Bauer Fostering Transdisciplinary Learning in Science and Technology	76
ESL7	Olgica Martinis The Current Challenges of Improving Chemistry Learning.....	77

POSTER PRESENTATIONS

P1	Wiktorija Adamska, Grzegorz Markiewicz, Artur R. Stefankiewicz Selective Guest Binding and Release: Tuning the Properties of Octameric Nanocapsules	79
P2	Marija Alešković, Amalija Ernečić, Sunčica Roca, Viktoriia Lazarenko, Nikola Bregović, Marina Šekutor Structural and Thermodynamic Features of Diamondoid Ammonium Salt Inclusion Complexes with Cyclodextrins	80
P3	Jasna Alić Stolar, Tomislav Stolar, Zoran Štefanić, Krunoslav Užarević, Marina Šekutor Sustainable Solution for Synthesis of Diamondoid Ethers.....	81
P4	Tea Babić, Andrea Usenik, Katarina Pičuljan, Nikola Bedeković, Josip Požar Complexation Thermodynamics of Benzene Derivatives with Macrocyclic Receptors	82
P5	Nikola Bedeković, Vladimir Stilinović, Dominik Cinčić Stoichiometric Diversity of Halogen Bonded Cocrystals Derived from <i>o</i> -Diiodotetrafluorobenzene and Monotopic Acceptors	83
P6	Branimir Bertoša, Zoe Jelić Matošević Molecular Modelling of MntR Transcription Factor from <i>Bacillus Subtilis</i>	84
P7	Katijana Bilandžić, Renata Odžak, Matilda Šprung Biological Activity of 3-Aminoquinuclidine Quarternary Salt	85
P8	Nataša Burić, Marina Šekutor Preparation and Mass Spectral Characterization of Diamondoid Derivatives	86
P9	Marija Cvetnić, Davor Mendeš, Nikola Bregović, Vladislav Tomišić Thermodynamics of Several Ion-Pairings in Acetonitrile.....	87
P10	Karla Dubaić, Leo Štefan, Maša Safundžić Kučuk Rheological Characterization – Important Tool in Reverse Engineering of Ophthalmic Solutions	88
P11	Ina Erceg, Marija Milić, Krunoslav Ilić, Darija Domazet Jurašin, Nadica Maltar Strmečki, Ivana Vinković Vrček, Maja Dutour Sikirić Calcium Phosphate/Titanate Nanomaterial Composites for Biomedical Applications.....	89
P12	Luka Fotović, Nikola Bedeković, Josip Požar, Vladimir Stilinović Can Halogen Bond Pay the Entropy Penalty?	90
P13	Josipa Golenja, Michael Sulyok, Ivana Dodlek Šarkanj, Bojan Šarkanj LC-MS/MS as a Solution in Precise, Selective, Confirmatory and Cost-Effective Food Multi-Contaminant Analysis	91
P14	Boris Gomaz, Zoran Štefanić Structure Comparison and Structural Alignments of Oligomeric Purine Nucleoside Phosphorylases	92

P15	<u>Gordan Horvat, Filip Kučas, Neven Golenić</u> A Computational Study of Calixarene-Cyclopeptide Anion Receptors	93
P16	<u>Suzana Inkret, Nadežda Radmilović, Kristina Smokrović, Nadica Maltar-Strmečki, Maja Dutour Sikirić</u> Influence of Differently Stabilized Silver Nanoparticles on Calcium Phosphate Formation and Transformation	94
P17	<u>Irena Ivanišević, Marko Zubak, Petar Kassal</u> Glass Surface Modification with Inkjet Printable Amphiphilic Silver Nanoparticles	95
P18	<u>Ivona Jasprica, Petar Horvat, Katarina Zrnc, Karl Bonney, Vidar Bjornstad, Ernest Meštrović, Robert Vianello, Lucija Hok, Nikola Bregović, Josip Požar, Katarina Leko, Vladislav Tomišić</u> Utilization of a Kinetic Isotope Effect to Decrease Decomposition of Ceftriaxone in a Mixture of D ₂ O/H ₂ O.....	96
P19	<u>Ilija Klarić, Jelena Kovačić, Mario-Livio Jeličić, Ana Mornar, Snježana Zubčić, Daniela Amidžić Klarić</u> Gluten in Dietary Supplements Used in the Treatment of Inflammatory Bowel Diseases.....	97
P20	<u>Jelena Kovačić, Mario-Livio Jeličić, Daniela Amidžić-Klarić, Ana Mornar</u> Optimization of LC-QqQ-MS/MS Method for Quantification of Andrographolides in Human Plasma	98
P21	<u>Željka Boček, Sara Krivačić, Petar Kassal</u> All-Solid-State Flexible Ammonium-Selective Electrode Based on Inkjet Printed Graphene Solid Contact	99
P22	<u>Marija Kuštro, Dubravka Tavra, Kristina Bule, Viktorija Prevarić, Matija Cvetnić, Šime Ukić, Tomislav Bolanča, Martina Miloloža, Marinko Markić, Lidija Furač, Dajana Kučić Grgić</u> Determination of the Optimal Conditions for the Biodegradation of Polyvinly Chloride by the Yeast <i>Candida Parapsilosis</i> Using a Full Factorial Plan	100
P23	<u>Dubravka Madunić-Čačić, Blaženka Čiško-Anić, Nikola Sakač, Marija Jozanović</u> Potentiometric Sensors and Water Solutions – Solution for Surfactant Analysis	101
P24	<u>Antonio Magnabosco, Gordana Majstorović, Tomislav Portada</u> Synthesis and Solvatochromism of Three Novel Brooker's Merocyanine Analogues	102
P25	<u>Lucija Marić, Nikola Bedeković, Vladimir Stilinović</u> Temperature Effect on Coordinative, Halogen and Tetrel Bonds in Crystal Structure Obtained from Lead Thiocyanate and 3-Iodopyridine	103
P26	<u>Marina Mavrinac, Vlasta Bertone</u> Impurities in Latanoprost Eye Drops: Analytical Procedure Lifecycle Management and QbD Approach in That Process	104
P27	<u>Mia Mesić, Tin Klačić, Davor Kovačević</u> Divalent Cation Effects on Poly(Diallyldimethylammonium Chloride)/Poly(Sodium 4-Styrenesulfonate) Nanofilm Deposition	105
P28	<u>Valentina Milašinić, Krešimir Molčanov, Petar Štrbac</u> Charge Density of Tetracyanoethylene (TCNE) Radical Anion	106
P29	<u>Matija Modrušan, Lucija Otmačić, Nikolina Vidović, Nikola Cindro, Ivo Crnolatac, Giovanna Speranza, Gordan Horvat, Vladislav Tomišić</u> Thermodynamics of Anion Binding by Pentaphenylalanine in Acetonitrile	107
P30	<u>Filip Perlitius, Anna Walczak, Miroslava Čonková, Grzegorz Markiewicz, Jack Harrowfield, Artur R. Stefankiewicz</u> Dimeric Capsule vs. Columnar Polymer: Structural Factors Determining the Solution and Solid-State Aggregation Behavior of Amino Acid Functionalized BTA Derivatives.....	108
P31	<u>Mateja Peršić, Mirela Samardžić, Andrea Dandić, Mateja Budetić</u> A New Solid-State Promethazine Hydrochloride-Selective Sensor Based on Functionalized MWCNT	109

P32	Martina Rajić, Mario-Livio Jeličić, Jelena Kovačić, Daniela Amidžić Klarić, Nikola Cindro, Ana Mornar Turk Characterization of Sulfonylureas' ADME Properties Using Biomimetic Chromatography and Computational Procedures	110
P33	Riya Sailani, Asha Rolaniya Ruthenium(III) Catalyzed Oxidative Degradation of Paracetamol by Hexacyanoferrate(III) in Solution Media - A Computational Screening and Mechanistic Pathway	111
P34	Filip Vranješević, Nikola Sakač, Marija Jozanović, Maria Kolypadi Markovic, Valerije Vrček, Dean Marković Cyclic Carbamate Formation through Palladium-Catalysed CO ₂ Absorption by Propargylic Amines	112
P35	Vesna Saršon, Josipa Šumonja Selection of the Delivery System for the Preservative Free Ophthalmic Product.....	113
P36	Sabino Sepčić, Patrizia Janković, Marko Babić, Daniela Kalafatović Design, Synthesis and Characterization of Linear and Cyclic Peptides	114
P37	Giuseppe Cappelletti, Eleonora Pargoletti, Marco Rabuffetti, Marina Simona Robescu, Sara Sangiorgio, Riccardo Semproli, Daniela Ubiali, Giovanna Speranza The BioSurf Project: A Sustainable Production of Bio-based Surfactants from Renewable Resources.....	115
P38	Petra Stanić, Krešimir Molčanov Stacking Patterns of Organic Radicals in Tetracyanoquinodimethane Salts	116
P39	Lu Turković, Zvonimir Mlinarić, Kristina Javorić, Juraj Vuić, Anamarija Skenderović, Sandro Makarić, Vladimir Radić, Tajana Silovski, Dragana Mutavdžić Pavlović, Miranda Sertić Optimization Studies of Solid Phase Extraction and LC-MS Analysis of CDK4/6 Inhibitors in Breast Cancer Combination Therapy in Plasma Samples	117
P40	Dorina Vidović, Nada Košutić Hulita, Nikola Marjanović Novel Techniques for Secondary Structure Determination of Peptides in Drug Development.....	118
P41	Sanja Vukoja, Sandra Siročić, Jelena Ranić, Tina Medančić, Diana Žgela, Lucija Hok, Ivona Jasprica, Robert Vianello, Ernest Meštrović Deep Eutectic Solvent System Based on Amino Acids Derivatives	119
P42	Jyoti Yadav, Thomas Kendall, Carmen Guguta, Swati A. Patel Nucleation Kinetics of Ascorbic Acid in Water and Water-Ethanol Solvent	120
P43	Marko Zubak, Sara Krivačić, Petar Kassal Photoreduction in Solutions by Intense Pulsed Light	121
P44	Kristina Đambić, Katarina Lubina, Dora Tot, Nikola Sakač, Dubravka Madunić-Čačić, Dean Marković, Bojan Šarkanj, Iva Pukleš, Nada Glumac, Marija Jozanović Synthesis of New Cationic Amphiphiles with Antimicrobial Activities	122
P45	Dario Matulja, Gabrijela Matijević, Sanja Babić, Maria Kolypadi Markovic, Marija Jozanović, Nikola Sakač, Rozelindra Čož-Rakovac, Sandra Kraljević Pavelić, Dean Marković Evaluation of Embryotoxicity of Granulatamide B and its Structural Analogues Using Zebrafish Model.....	123

AUTHORS AND PARTICIPANTS

AUTHORS	124
PARTICIPANTS	126

PLENARY LECTURES

SOLUTIONS IN SUPRAMOLECULAR CHEMISTRY IN WATER

Werner M. Nau

School of Science, Jacobs University Bremen, Campus Ring 1, 28759 Bremen, Germany

w.nau@jacobs-university.de

Supramolecular chemists have explored and developed a large portfolio of noncovalent driving forces to understand and ultimately control assembly formation. The formation of discrete host-guest complexes in aqueous solution has proven particularly instructive in this respect, also due to its implications for biological and environmental recognition events. Besides electrostatic interactions between host and guest, the driving force portfolio in water has been limited to two attractive interactions. These include the hydrophobic effect and dispersion interactions, which are additionally difficult to dissect. Here we present results which – conceptually – expand the portfolio of noncovalent driving forces in aqueous host-guest chemistry. First, besides the classical, entropically driven hydrophobic effect related to guest desolvation, a nonclassical, enthalpically driven counterpart related to the desolvation of intermediary sized host cavities needs to be considered, which can be traced back to the removal of high-energy cavity water as the driving force.¹ Second, the hydrophobic effect can be further dissected into two separate components, one related to differential cavitation energies, the other one related to differential dispersion interactions.² In supramolecular design, both need to be optimized separately, with the result that very small, non-solvated cavities contribute an extra driving force in the binding of very small guests such as gases; this can be exploited, for example, for gas-storage materials. Finally, through recent studies in the binding of large, so-called superchaotropic, anions to hydrophobic cavities, the chaotropic effect has emerged as a thermodynamically orthogonal driving force to the hydrophobic effect.³ The two effects derive from a different aqueous solvation pattern of hydrophobic and chaotropic guest molecules, which ultimately drives their desolvation and binding to hydrophobic cavities either entropically (hydrophobic) or enthalpically (chaotropic), see Fig. 1. Most recently, the first biological application of the chaotropic effect has been introduced, which related to the use of superchaotropic ions for transmembrane transport of amino acids, peptides, and drugs.⁴




Figure 1. Continuum of aqueous solvation patterns and associated driving forces.

REFERENCES

1. F. Biedermann, W. M. Nau, H.-J. Schneider, *Angew. Chem. Int. Ed.* **2014**, *53*, 11158.
2. S. He, F. Biedermann, N. Vankova, L. Zhechkov, T. Heine, R. E. Hoffman, A. De Simone, T. T. Duignan, W. M. Nau, *Nat. Chem.* **2018**, *10*, 1252.
3. K. I. Assaf, W. M. Nau, *Angew. Chem. Int. Ed.* **2018**, *57*, 13968.
4. A. Barba-Bon, G. Salluce, I. Lostalé-Seijo, K. I. Assaf, A. Hennig, J. Montenegro, W. M. Nau, *Nature* **2022**, *603*, 637.

SENSING SOLUTIONS WITH POTENTIOMETRIC PROBES

Eric Bakker,* Tara Forrest, Elena Zdrachek, Polyxeni Damala, Pitchnaree Kraikaew,
Robin Nussbaum

Department of Inorganic and Analytical Chemistry, University of Geneva, Quai E.-Ansermet 30,
1211 Geneva 4, Switzerland

* eric.bakker@unige.ch

Throughout the history of potentiometric sensors, the concept of symmetry has been central for assuring optimal signal stability. With pH electrodes, for example, the internal Ag/AgCl element at the indicator electrode side is accompanied with a similar Ag/AgCl electrode at the reference electrode side. The overall cell potential no longer depends on the specific redox couple because the two are compensated. Commercial pH probes of high E^0 potential reproducibility and stability are designed for the two redox elements to be similarly configured and placed within the sensor body so that temperature fluctuations affect both elements at very similar rates. This concept of symmetry was also used by Simon for the realization of a calibration-free potentiometric sensor where the observed value of the cell potential gives direct information about sample ion activity.¹

This symmetry is broken when the internal Ag/AgCl element of an indicator electrode is replaced with a solid contact transducing material because the reference element is now formed by a different redox couple. This is unfortunate because a major research effort of recent years has been the aim to realize highly stable all-solid-contact potentiometric probes, which is not helped by the asymmetric nature of the cell.

This talk will underline the importance of symmetry in such electrochemical sensors and show examples on how it can be restored with modern all-solid-state potentiometric probes. Examples will include a recently published example for the detection of the nutrient nitrate² where nitrate is added to the bridge electrolyte of the reference element so that a second nitrate electrode of the same design, rather than an Ag/AgCl can be used. This study will then be extended to solid contact pH probes that may behave in complete analogy to commercial pH probes, with a zero potential value at pH 7. Symmetry may also help to avoid the lengthy conditioning periods often required for best stability. Moreover, it is shown that when the reference element is matrix matched to the sample, the sensor can be made to directly output concentrations, rather than ion activities. The concept of symmetry is then applied to the design of submersible aquatic sensing probes, which are deployed successfully in freshwater systems in Switzerland.

REFERENCES

1. G. Rumpf, U. Spichiger-Keller, H. Bühler, W. Simon, *Anal. Sci.* 1992, 8, 553.
2. E. Zdrachek, T. Forrest, E. Bakker, *Anal. Chem.* 2021, 94, 612.
3. P. Damala, E. Zdrachek, T. Forrest, E. Bakker, *Anal. Chem.* 2022, 94, 11549.


Every Solution has a Question!

Prepare your team to participate in the **AlphaChrom pub quiz** during the Solutions in Chemistry conference.

Where, when, how... find out at the registration desk.

Don't worry, we have all the answers.

And prizes for you too!



THE HUMAN FACTOR IN SCIENTIFIC THINKING: THE ILLUSIONS THAT WE LIVE BY

Csaba Szántay

Spectroscopic Research Department, Gedeon Richter Plc. (www.richter.hu)

H-1475 Budapest 10., P.O.B. 27, HUNGARY

cs.szantay@richer.hu

The talk is based on a book published in 2015, and takes a candid look at science from an unusual perspective, attempting to offer some insights into how psychological factors secretly infiltrate its fabric.¹ The common notion that science is the world of objective rationality based on methodological rigor, stringent logic, irrefutable experimental proofs, and unbiased peer reviewing and testing, is an idealistic myth which all too often dissents from what “scientific truth” is in reality, and from how scientists truly „function” in practice. Actually, our scientific thinking is influenced by deep-rooted human (“anthropic”) factors of which we are not normally aware of. Some of these lead to what we call mental traps, i.e., the hidden sources of mistaken or misleading inferences, a phenomenon we refer to as the illusion of understanding, and mass misconceptions about apparently well-established “scientific truths”. By their very nature, mental traps affect even the smartest, most knowledgeable, and most attentive scientists. However, by understanding the essence of the traps one can develop the enlightening faculty of detecting and avoiding them both in one’s own and in others’ thoughts. It is this mental aptitude/attitude of being keenly conscious about our human nature during scientific thinking which is captured in the phrase “anthropic awareness” in the title of the aforementioned book. The talk will address the concept of “anthropic awareness” as a kind of philosophy, will outline the reasons behind the mental traps, and will discuss some of the traps, including a few real-life examples. I will attempt to show that “anthropic awareness” has a general relevance throughout all of natural sciences, and is useful not only in our everyday professional lives as researchers, but also in our nonprofessional everyday lives.

REFERENCES

1. Cs. Szántay, Jr, (Ed), *Anthropic Awareness: The Human Aspects of Scientific Thinking in NMR Spectroscopy and Mass Spectrometry*. New York: Elsevier, 2015.

ADVENTURES IN PHASE SPACE OF DNA BINDING AND STABILITY

Jurij Lah,* San Hadži

University of Ljubljana, Faculty of Chemistry and Chemical Technology, Večna pot,
1000 Ljubljana, Slovenia

* jurij.lah@fkkt.uni-lj.si

In 1953, Watson and Crick presented their iconic structure of double-stranded DNA, which has become firmly imprinted in the cultural memory of mankind. However, DNA is capable of adopting other structures that occur in the context of regulating common and disease-related cellular processes, leading to complex DNA stability landscapes. Here we show how thermodynamics can be used to estimate the mechanism of DNA folding and structural transformation processes, determining their driving forces, and for predicting DNA transcription regulation properties. This will be demonstrated using two model systems:

(i) Guanine-rich DNA can form G-quadruplex structures in key regions of the genome, including promoter regions, oncogenes, and telomeres. We will address one of the key questions in G-quadruplex formation, namely why a single guanine-rich sequence can fold into and switch between different G-quadruplex structures depending on temperature and the presence of co-solvents, co-solutes and ligands.¹⁻⁴

(ii) The bacterial stress response through genetic systems called toxin-antitoxin modules (TA) could be a major cause of chronic infections. TA modules encode a stable toxin (T) and an unstable antitoxin (A) responsible for the formation of the transcription repressor complexes with DNA. We will explain the molecular mechanism of transcriptional regulation of the TA module.⁵

The thermodynamic characterization of these systems is based on the global analysis of a variety of calorimetry (DSC, ITC) and spectroscopy (CD, fluorescence) data [1-3]. We will show how the obtained thermodynamic parameters can be related to structural properties and used to predict DNA behavior under different conditions.

ACKNOWLEDGEMENTS The financial support of the Slovenian Research Agency projects P1-0201 and J1-1706 is gratefully acknowledged.

REFERENCES

1. S. Hadži, V. Kocman, D. Oblak., J. Plavec, J. Lah, *Angew. Chem. Int. Ed.* 2019, *58*, 2387.
2. M. Bončina, G. Vesnaver, J. B. Chaires, J. Lah, *Angew. Chem. Int. Ed.* 2016, *55*, 10340.
3. D. Oblak, S. Hadži, Č. Podlipnik, J. Lah, *Pharmaceuticals* 2022, *15*, 1.
4. M. Bončina, et al., and J. Lah, *Nucleic Acids Res.* 2015, *43*, 10376.
5. A. Vandervelde, et al. and J. Lah, *Nucleic Acids Res.* 2017, *45*, 2937.

JGL Establishes Scientific Advisory Board to Advance the Development of Innovative and Complex Products



This is a big step forward for us, and for Croatia as a whole, because we are developing science-based, advanced products tailored to the needs of patients and consumers, says JGL

The JGL Scientific Advisory Board was established in September 2021 with the aim of positioning JGL as an innovator in the development of advanced, differentiated products in three strategic therapeutic areas – flu and cold, ophthalmology and dermatology. This independent body comprises top domestic and foreign experts and raises the science profile of the company's R&D projects. The company now operates in 60 markets around the world, either directly or through partners, and its portfolio consists of 150 brands, 300 products and 650 variations tailored to the needs of specific markets.

- The establishment of the Scientific Advisory Board is another step forward in the development of innovative products around which we build our science and innovation ecosystem. The goal of this body is to cooperate with the JGL team and steer existing and develop new ideas for the company's manufacturing and technology portfolio, all of it based on sound science, says Mislav Vučić, CEO of JGL. He also adds that he is very proud of the combination of cutting-edge science and talent, especially during their annual in-person meetings, where they network and share relevant, science-based knowledge and ideas through synergy and lay the groundwork for increasing the company's competitiveness.

Second SAB meeting was organized in September 2022. The Advisory Board consists of nine experts, internationally recognised in their respective fields,

same being of high relevance for JGL projects. They include Prof. Henning H. Blume, PhD, Prof. Ralph Mösges, PhD, Prof. Stipan Jonjić, PhD, Jag Ahluwalia, PhD, Prof. Özgen Özer, PhD, Prof. Vladimir Trkulja, PhD, Chief Physician Sonja Jandroković, PhD, Andreas Bilstein, PhD, and Prof. Jasmina Lovrić, PhD. The team is coordinated by Zdravka Knežević, PhD, Director of Scientific Operations at JGL, who participated in a number of similar initiatives and projects and has over 30 years of experience in R&D, drug registration, and clinical development.

- I am delighted that I get to share my knowledge and experience from international companies at JGL, as well as cooperate with a great team of experts from the EU, UK, Turkey and other global markets on the Board. I believe that this represents great progress for JGL and our teams, as well as for Croatia as a whole, because we will be able to develop new products for the benefit of our patients and the treatment needs of modern society, with clear scientific rationale, emphasises Knežević.

In cooperation with SAB Memebres, relevant JGL teams conducted a series of innovative formulation, preclinical, clinical, and analytical studies in 2021 and 2022, all with the aim of creating a network of relevant knowledge and identifying new projects for the company's therapeutic areas and technology portfolio.





The JGL Scientific Advisory Board, with Zdravka Knežević, PhD, Director of Scientific Operations

A Major European Manufacturer of Sterile Forms

For more than 30 years, JGL has been growing and developing its business globally. Thanks to continued investments in R&D, production facilities and technology, it currently ranks among the leading manufacturers of sterile pharmaceutical forms in the EU. The business, research and manufacturing complex JGL Pharma Valley in Rijeka combines best practices in production process management, state-of-the-art technology in pharmaceutical production and high environmental standards.

JGL is currently implementing a new investment project, Integra 2020, worth HRK 375m, which will integrate the company's development, production, and storage capacities and encompass three units – R&D, commercial production and logistics centre.

Nearly 85 Per Cent of Revenue Comes From Exports

The JGL Group ended 2021 with total revenue of HRK 1.1 billion and a profit before tax of more than HRK 100 million. Nearly 85 per cent of the company's core earnings is generated outside of Croatia, and JGL Group employs just over a thousand people.

The company's leading export brand is Aqua Maris, launched in 1999, which consists of 100% natural products based on seawater from the Adriatic, used for the prevention and treatment of upper respiratory tract diseases. Other key brands include Vizol S, Meralys, Aknekutan and Dramina.



Prof. Ralph Mösges, PhD, a world-renowned expert in allergology, pulmonology and clinical immunology



Prof. Henning H. Blume, PhD, who specialises in biopharmaceutical and pharmacokinetic research, with a particular focus on characterising the bioavailability and bioequivalence of drugs

SUPRAMOLECULAR CHEMISTRY: SOLUTION THERMODYNAMICS AND ITS APPLICATIONS

Angela F. Danil de Namor

Department of Chemistry, University of Surrey, Guildford, Surrey GU2 7ST

A.Danil-De-Namor@surrey.ac.uk

Following a brief introduction on the strategy adopted in our research on the thermodynamics of Supramolecular Chemistry involving neutral and ionic species and their applications, the lecture will address some aspects related to the thermodynamics of common salts in low permittivity media and their use in several processes. Given the scope of Supramolecular Chemistry for the availability of new salts in which one of the components is a complexed anion or cation resulting from the ion- receptor interaction, advances so far made by our Group on their behaviour in solution will be based on research carried out with a wide range of macrocycles such as crown ethers, cryptands¹ and calix [4] based macrocyclic receptors.^{2–4} Evidence supporting the solvation changes observed in the reactants and the product resulting from their transfer from one solvent to another will be given. Their contribution in assessing the main factors controlling the selective, non-selective or non-interactive behaviour of the receptor towards the guest in solution processes as well as the relevance of solution thermodynamics in the design of new materials for various purposes will be illustrated with representative examples.⁵

ACKNOWLEDGEMENTS A F D de N thanks the European Commission for the financial support given under various Framework Programmes, the Science & Engineering Research Council (UK) and the Leverhulme Trust Foundation for financial support.

REFERENCES

1. R M Izatt, K Pawlak, J S Bradshaw, *Chem. Rev.* **1991**, *91*, 1721.
2. C D Gutsche, Chapter 1 in *Calixarenes 2001*. Eds Z Asfari, V Böhmer, J Harrowfield, J Vicens, Kluwer Academic Publishers, **2001**.
3. A F Danil de Namor, Chapter 19 in *Calixarenes 2001*. Eds Z Asfari, V Böhmer, J Harrowfield, J Vicens, Kluwer Academic Publishers, **2001**.
4. P A Gale, P Anzenbacher, J L Sessler, *Coord. Chem. Rev.* **2001**, *222*, 57.
5. A F Danil de Namor, *DSc Thesis*, University of Surrey, **2022** and references within

CHEMICAL SOLUTIONS FOR AGRI-FOOD WASTE UPCYCLING

Nicoletta Ravasio

CNR SCITEC, via C.Golgi 19, 20133 Milano, Italy

nicoletta.ravasio@scitec.cnr.it

The food supply chain generates enormous amount of waste: veggies residues left over in the fields, side stream of food production, post-consumer waste. Most of them are used as feed or in bioenergy production, some are disposed of as waste.

However, a green chemistry approach can allow one, once the different chemical classes present in the waste are identified, to design specific valorization pathways for each of them also outside the food value chain. Two examples will be reported and discussed: the valorization of rice waste carried out in the frame of RiceRes project and the upcycling of silverskin, a waste of coffee roasting, in the CirCo project.

The multivalORIZATION of rice waste led to green building materials, innovative polymer matrix composites, sterolesters for nutraceutical use and high added value protein hydrolysates.¹ Both chemical and enzymatic processes were used for the upgrade of the cellulose, the lignin, the oil and the protein fraction of straw, husk and bran.

Extraction of silverskin with supercritical CO₂ led to a fat of unusual composition that was found to be suitable for the formulation of makeup products.² The subsequent extraction of a powerful antioxidant and its esterification by heterogeneous catalysis led to a product with increased antioxidant properties. Finally, the residue was used as a replacement of virgin cellulose in the production of graphic paper. A detailed LCA analysis allowed to state that substitution of 15 % virgin cellulose with silverskin derived cellulose can reduce the environmental impact of paper production by 10 % and greenhouse gas (GHG) emissions by 13 % compared to conventional production.³

The hydrogenation of lactose, the main component of cheese whey, to sorbitol and dulcitol will also be discussed⁴.

REFERENCES

1. E. Overturf, N. Ravasio, F. Zaccheria, C. Tonin, A. Patrucco, F. Bertini, M. Canetti, K. Avramidou, G. Speranza, T. Bavaro, D. Ubiali, *Bioresource Technology Reports* **2020**, *11*, 100427.
2. R. Nasti, A. Galeazzi, S. Marzorati, F. Zaccheria, N. Ravasio, G.L.Bozzano, F. Manenti, L. Verotta, *Waste and Biomass Valorization*, **2021**, *12*, 6021.
3. E. Overturf, S. Pezzutto, M. Boschiero, N. Ravasio, A. Monegato, *Sustainability* **2021**, *13*, 9069.
4. F. Zaccheria, M. Mariani, N. Scotti, R.Psaro, N. Ravasio, *Green Chemistry*, **2017**, *19*, 1904.

***“Invaluable instrument
in pharmaceutical
process development”***

*Principal Scientist MSD,
Lead of Crystallization Lab.*



**DESIGNED BY SCIENTISTS
FOR SCIENTISTS**

**Crystal
16**

A POWERFUL SYSTEM FOR POLYMORPH, SALT & CO-CRYSTALLIZATION RESEARCH

- 4 solubility curves in 4 hours
- Obtain both solubility curve and MSZW information simultaneously
- Design quickly any crystallization process
- Screen for solvents or any solid-state form

Please contact us for more information

info@crystallizationsystems.com

crystallizationsystems.com

Tel: +31 72 30 200 40

Pyrietstraat 2

1812 SC Alkmaar, The Netherlands

Follow us on  


Technobis

CRYSTALLIZATION SYSTEMS


DISCOVER SCREEN OPTIMIZE

You are now able to perform well controlled crystallization studies from hit identification and lead identification up through process scale up. Combine the *CrystalBreeder*, *Crystal16* and *Crystalline* in a flexible configuration to optimize solid-state success!


Single Crystal Growth



Solubility curves in different solvents



Seeding Crystallization



Discover

Discover new polymorphs or grow single crystals with success using different controlled crystallization methods in parallel such as:

- thermal evaporation
- solvent evaporation
- vapor diffusion
- sublimation



Crystal
BREEDER

Screen

Screen for selecting solvents, polymorphs, salts or co-crystals has never been so easy. The world's most used equipment for solubility curves measurement and phase diagrams.



Crystal
16

Optimize

Optimize your crystallization or formulation process at an early stage with the ultimate particle view imaging technology and Raman Spectroscopy.



Crystal
LINE

Please contact us for more information

info@crystallizationsystems.com

crystallizationsystems.com

Tel: +31 72 30 200 40

Pyrietstraat 2

1812 SC Alkmaar, The Netherlands

Follow us on  

Technobis
CRYSTALLIZATION SYSTEMS

INVITED LECTURES

POTENTIAL SOLUTION FOR PEPTIDE-BASED DRUG DELIVERY TO THE BRAIN: DESIGN AND ACTIVITY OF MULTIFUNCTIONAL CYCLIC PEPTIDE-BASED ANALGESICS

Predrag Cudic,^{a,*} Shahayra Majumder,^b Heather Stacy^b, Thomas Cirino,^b Brandon Williams,^a Jay P. McLaughlin^b

^a Department of Chemistry and Biochemistry, Charles E. Schmidt College of Science, Florida Atlantic University, Boca Raton, FL 3343, USA, ^b Department of Pharmacodynamics, College of Pharmacy, University of Florida, Gainesville, FL 32610, USA

* pcudic@fau.edu

Chronic pain is the leading cause of disability worldwide, accounting for significant hospitalization, prolonged care and economic cost. Most clinically-used opioid drugs are μ -opioid receptor (MOR) agonists with liabilities of tolerance, high potential for addiction and respiratory depression mediated by this receptor.¹ In contrast, κ -Opioid receptor (KOR) agonists are effective pain suppressors and unlike MOR agonists do not cause respiratory failure or addiction¹. However, KOR agonists developed to date exhibit undesirable dysphoria and aversion attributed to KOR suppression of dopamine and serotonin. Given the severity of the problem, there is an undeniable and urgent need for new pain medication without the unwanted side-effects of existing opioid drugs. Among the chemical entities utilized in drug discovery, peptides emerge as particularly attractive lead compounds for the discovery and development of new central nervous system (CNS) drugs due to their roles as potent regulators of normal CNS function. However, conventional drug delivery methods (oral and intravenous) are inefficient in delivering hydrophilic and high molecular weight drugs, such as peptides and proteins to the brain. Both the blood-brain barrier (BBB) and blood-cerebrospinal fluid barrier (BCB) restrict the transport of these therapeutic agents from systemic circulation into the CNS. Alternatively, intranasal (i.n.) administration may enable these therapeutic agents to directly enter the brain by bypassing the BBB and BCB, avoiding systemic circulation of the drug and reducing the risk of systemic side effects as well as hepatic/renal clearing and toxicity. To address the need for novel therapeutics to treat chronic pain, we designed new cyclic peptide-based analgesics suitable for i.n. delivery that combine selective KOR agonist and antidepressant pain relieving mechanisms. This is achieved by grafting bioactive analgesic sequences into a carrier cyclic peptide scaffold exhibiting bioadhesive properties.^{2,3} Combining KOR agonist activity with simultaneous delivery of serotonin may improve chronic pain management when more than one physiological mechanism or system is implicated and may potentially minimize side effects associated with the KOR activation.

ACKNOWLEDGEMENTS: This work was supported by the NIH (NIDA) RDA039722A grant to P.C.

REFERENCES

1. R. Przewlocki, B. Przewlocka, *Eur. J. Pharm.* 2001; 429, 79.
2. Y. Singh, P. Cudic, M. Cudic, *Eur. J. Org. Chem.* 2022, 28, e202200302.
3. M. Rodriguez, A. B. Yongye, M. Cudic, K. M. Mayorga, *et al.*, *Amino Acids*, 2017, 49, 1867.

NMR SPECTROSCOPY AND GENERIC PHARMA R&D: PAST, PRESENT AND THE FUTURE

Tomislav Biljan

Global R&D, Pliva Croatia Ltd., Prilaz baruna Filipovića 29, Zagreb, Croatia

tomislav.biljan@pliva.hr

Past, present and future use of NMR spectroscopy in generic pharma industry is presented. Special emphasis is given to the following aspects relevant for generic pharma industry: organic synthesis, kinetics, impurities, catalysts, quantitative NMR, beyond proton and carbon, solid state NMR, polymorphs, excipients, drug products, quality and regulatory aspects. These aspects are highlighted by the use of real life examples.



Xellia Pharmaceuticals is a specialty pharmaceutical company focused on providing important anti-infective treatments against serious and often life-threatening infections. Our anti-infective treatments are generics that combat serious bacterial and antibiotic resistant infections as well as certain fungal diseases. We are a world leader in the development, manufacturing and supply of fermented anti-infectives sold as active pharmaceutical ingredients and finished dosage forms to key pharmaceutical industry companies. Headquartered in Copenhagen, Denmark and owned by Novo Holdings A/S, Xellia Pharmaceuticals has more than 1500 employees globally. From state-of-the-art manufacturing sites in the U.S., China, Denmark and Hungary to R&D sites in Norway and Croatia; Xellia Pharmaceuticals excels within innovative product development to deliver high quality products to its customers.

Our product portfolio of anti-bacterial and anti-fungal products used for the treatment of severe infections, including infections caused by multi-resistant bacteria and fungi, is the foundation for our growth strategy within the antibiotics sector. Xellia is the leading supplier of important anti-infectives Vancomycin and Colistimethate Sodium (CMS). Our success and strong market position is built on more than 100 years experience in the pharmaceutical industry. Xellia Pharmaceuticals' focus its R&D investments within inhalable and injectable product technologies as we are committed to researching solutions, which not only improve patients' quality of life, but will also save lives. Based on our long-term expertise, we bring value for our stakeholders and signify a patient-centric mindset through an embedded corporate motivation of integrating science and innovation to save lives.



MOLECULES BEHIND WINE TASTING

Luna Maslov Bandić

University of Zagreb, Faculty of Agriculture, Svetošimunska c. 25, 10000 Zagreb, Croatia

lmaslov@agr.hr

Wine has played and continues to play a role in a variety of religious, cultural, and social activities. Despite its almost seven millennia as a part of the human experience, our comprehension of this beverage and all of its facts remains woefully lacking, especially in light of modern science's capabilities. Wine is a fascinating chemical mixture. Each bottle contains thousands, if not tens of thousands, of different molecules ranging from acids and sugars to phenolic compounds and vanishingly low concentration aroma compounds. Although wine is sometimes offered with an impression of ancient heritage, industry uses modern techniques and analysis. There is an increasing number of examples where flavor molecules have been found and winemakers have used this information to improve their product. The most common methoxypyrazine found in wine, 2-isobutyl-3-methoxypyrazine (IBMP), provides a distinctive green bell pepper scent to particular wine grape varieties (i.e. Sauvignon blanc, Cabernet Sauvignon, etc.) with a sensory threshold ranging from 8 to 15 ng L⁻¹ depending on the wine matrix¹. Methoxypyrazines are of special importance in the wine business; while some varieties of wine are recognized for high methoxypyrazine levels (e.g., New Zealand style Sauvignon blanc), consumers may reject wines even when methoxypyrazine concentrations are in the low ng L⁻¹ range. One of the most challenging aspects of methoxypyrazine analysis in wine is achieving the required analytical sensitivity. Rotundone, a sesquiterpene, was discovered as the main contributor to the aroma of black pepper in wine in 2008. This powerful flavor component has now been discovered in other grape types. Rotundone is generally well received by consumers. One of the most remarkable facts about this chemical is that it is anosmic to 20 % to 25% of the general population. According to a research of Italian *V. vinifera* varieties, furaneol is important to the aroma of Refosco and Primitivo (also known as Zinfandel in the United States and Crljenak kaštelski or Tribidrag in Croatia). Furaneol and homofuraneol both have low odor thresholds, 5 and 125 µg/L, and have an additive and/or synergistic role in expressing the fruity and caramel flavor of rosé wines². It is clear that wine's aroma depends not on a single compound but rather on the composition and interactions of the many odor-active compounds present. The potential aroma of wine is also dependent on the release of aroma compounds from their odorless precursors during wine maturation and the modification of volatiles due to chemical changes. Future discoveries will continue to be driven by development of improved and high-throughput analytical methods that will allow monitoring of a large number of volatiles, including those present at low concentrations.

REFERENCES

1. I. Ryona, S. Leclerc, G. L. Sacks, *Journal of Agricultural and Food chemistry* **2010**, *58*, 9723.
2. A. L. Robinson, P. K. Boss, P. S. Solomon, R. D. Trenogove, H. Heymann, S. E. Ebeler, *American Journal of Viticulture and Enology* **2014**, *65*, 1.

SOLVING COMPLEX PROBLEMS IN SOLUTION CHEMISTRY BY DESIGN OF EXPERIMENT

Iva Rezić

Department of Applied Chemistry, University of Zagreb Faculty of Textile Technology,
Prilaz baruna Filipovića 28a 10000 Zagreb, Croatia

iva_rezic@net.hr, iva.rezic@ttf.hr

In process and product development, e.g. formulation of new products, the solvent system is under control of many stochastic parameters so it cannot be fully described or optimized without the sophisticated statistical models and tools.

This presentation will therefore give an overview of several examples, from laboratory to industrial scale, in which the complex systems were optimized by using the statistical methodology *Design of Experiment (DoE)*. The examples will cover:

- i) the formulation of new products (detergents) [1],
- ii) the development of new method (chromatographic separation of amino-acids and ultrasonic extraction) [2,3] and
- iii) the optimization of the industrial process for achieving the best product performance (maximal hydrophobic coating) [4].

The results of the investigations showed that the DoE is an effective tool that significantly enabled reducing time and money required for the formulation of a new product, or optimization and manipulation with the new processes (technological, industrial and scientific). Moreover, the estimation of the influence and interactions between all factors influencing the systems was enabled. The biggest advantage of a DOE analysis was that it provided solutions that can lead to faster time to market, lower development costs, lower operating costs, and lower cost of poor product quality. Thus, DoE is an effective and unavoidable tool for chemists dealing with complex solvent systems.

ACKNOWLEDGEMENTS This research was partially funded by the Croatian Science Foundation grant number IP-2019-04-1381 (project under the title 'Antibacterial coating for biodegradable medicine materials ABBAMEDICA'). Any opinions, findings, conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the Croatian Science Foundation.

REFERENCES

1. I. Rezić, *Monatshefte für Chemie –Chemical Monthly* 2011, 142, 1219.
2. I. Rezić et al. *Journal of Planar Chromatography Modern TLC* 2007, 20, 173.
3. I. Rezić, *Ultrasonic Sonochemistry*, 2009, 16, 63.
4. I. Rezić and A. Kiš, *Polymers*, 2020, 12, 2131.



EuChemS

European Chemical Society

EuChemS, the European Chemical Society, aims to nurture a platform for scientific discussion and to provide a single, unbiased European voice on key policy issues in chemistry and related fields.

Representing more than 160,000 chemists from more than 40 Member Societies and other chemistry related organisations, EuChemS relies on a unique network of active researchers involved in all the fields of chemistry. Through this network, EuChemS organises several specialised academic conferences as well as the biannual EuChemS Chemistry Congress, the European congress of chemical sciences. EuChemS also promotes the role and image of the chemical sciences among the general public and policy-makers through social media, newsletters and through the organisation of conferences and workshops open to the society.

Through the promotion of chemistry and by providing expert and scientific advice, EuChemS aims to take part of the solution to today's major societal challenges.

For more information about the European Chemical Society (EuChemS), please visit www.euchems.eu or contact us at:

EuChemS aisbl
Rue du Trône 62
1050 - Brussels
Belgium

Phone: +32 2289 25 67 | +32 2289 26 90

Email: secretariat@euchems.eu

 <http://on.fb.me/1B8Qa0n>

 <https://twitter.com/EuCheMS>

THE PATH TOWARDS ENLIGHTENMENT? SUPPORTING STUDENTS IN SOLVING COMPLEX CHEMICAL CALCULATIONS USING AN ADAPTIVE DIGITAL PLATFORM

Ewald Edink,^{a,*} Jeroen Bottema,^b Artur Lorincz,^c Kinga Lorincz^c

^a Department of Life Sciences & Chemistry, Faculty of Agri, Food & Life Sciences, Inholland University of applied sciences, De Boelelaan 1109, 1081 HV Amsterdam, The Netherlands, ^b Department of Teaching, Learning & Technology, Inholland University of applied sciences, Theresiastraat 8, 2593 AN Den Haag, The Netherlands, ^c Enlight Ed, Fred. Roeskestraat 115, 1076 EE Amsterdam, The Netherlands

* ewald.edink@inholland.nl

A large number of our undergraduate students at the Inholland University department of Life Sciences & Chemistry encounter difficulties and frustrations when learning to solve (complex) chemical quantitative problems. This is problematic since solving chemical calculations is considered a crucial skill for the profession for which we train our students, i.e. B.Sc. level laboratory analysts. In order to identify the most common reasons why students have difficulties with learning this important skill, a literature survey has been performed. This survey resulted in identification of a large set of articles and studies addressing this specific topic, exemplifying that this is a widespread issue in secondary and higher educational settings.¹⁻⁴

A possible solution is provided by Enlight-Ed's online digital adaptive platform⁵ that enables simulation of the conversation between learner and teacher in a classroom when the learner is having difficulties solving assignments. Using a chatbot functionality, students are exposed to adaptive assignments that resemble the 1-to-1 learning experience and optimise student engagement. The platform is also capable of collecting learning data providing the teacher an overview on the level of students and common misconceptions allowing to make informed decisions.

Using the editing functionality of the Enlight-Ed platform and the obtained insights from the aforementioned literature survey, adaptive assignments were specifically created for our first-year Life Sciences & Chemistry students. These digital interactive assignments assist students in following the step-wise approach that is needed to solve such complex assignments. When needed, the students is provided with additional instructions when needed. An evaluation poll amongst participants of this pilot-study showed high appreciation of this learning methodology. 77% of respondents agreed that the adaptive assignments using the Enlight Ed platform helped them in understanding how to solve such complex chemical calculations.⁶

REFERENCES

1. C. Furio, R. Azcona, G Jenaro, *Chem. Educ. Res. Pract.* 2002, 3, 277.
2. G. M. Bodner, *Chemistry education*. 2015, Weinheim, Germany: Wiley-VCH Verlag GmbH & Co, 181.
3. K. Salta, C. Tzougraki, *Research in Science Education* 2010, 41, 587.
4. J. Staver, A. Lumpe, *Journal of Research in Science Teaching* 1995, 32, 177.
5. <https://enlighted.app/>
6. E. Edink, <https://media-and-learning.eu/type/featured-articles/the-path-towards-enlightment-supporting-students-in-complex-problem-solving-using-enlight-ed/>

CHEMISTRY IN THREE DIMENSIONS: CHIRALITY, SYMMETRY, ISOMERISM

Krešimir Molčanov

Department of Physical Chemistry, Ruđer Bošković Institute, Bijenička c. 54, HR-10000 Zagreb, Croatia

kmolcano@irb.hr

Poor understanding of fundamental concepts of stereochemistry, such as enantiomerism or stereoisomerism, is unfortunately rather common among Croatian chemists and chemistry teachers. The reason for this is an awkward approach to education, which is based on erroneous foundations. The basic stereochemical concepts in Croatia are introduced through definition of constitutional isomerism and stereoisomerism, and definition of the "asymmetric" carbon atom. Such an approach yields questionable results: according to the author's experience with students of chemistry, graduated chemists, chemistry teachers and even some researchers, the majority of Croatian chemists actually don't understand what symmetry and chirality are.

It is well known that atoms and molecules are 3D objects, and their representation in two dimensions (on a blackboard, paper or screen) is merely projection. Therefore, for understanding basic chemical concepts which involve three dimensions - chirality, isomerism and symmetry - 3D models are indispensable. Fortunately, dedicated molecular model kits (which are expensive) are not needed. Many objects of every-day use can find a new use as models for teaching stereochemical concepts, such as symmetry (flowers, building blocks, etc.) or chirality (screws, shoes, corkscrews, snails). Toy building blocks can be used to demonstrate concepts of constitutional isomerism and stereoisomerism and also to assemble simple chiral objects.

A chiral object can be converted into its mirror image simply by watching it in a mirror. By observation of common things around us we can understand the phenomena of chirality and enantiomerism; then we make a mathematical abstraction - i.e. how to define a helix or a left or a right coordinate system, etc. Molecular chirality then becomes an easily understood phenomenon.

Thus, in teaching chemistry we can dispense with "chirality centres" and *R* and *S* configurations; they are not just needless, but superfluous. Tetrahedral carbon atom of van't Hoff and Le Bel is of a great historical importance, but it is just a special case of a common phenomenon of molecular chirality: "chirality centre" is not necessarily a tetrahedral carbon atom, it can be any atom with 4 or more atoms bound to it; there are also chiral molecules lacking "chiral centres", and they may possess quite a high symmetry. Instead of insisting on conventions and bookkeeping devices (such as *R* and *S* configurations and similar IUPAC rules), we should first focus on the basic concepts and introduce them through every-day objects. After that we can move to a more abstract, "mathematical" definition; conventions should only come at the end.

LEUKOCARE

Biopharmaceutical Drug Product Development Experts



Choose the leading innovator
in best-in-class biopharmaceutical drug product development
to support your development of novel therapies

Best formulation for your drug product

Broader Design Space

Only with Leukocare you will have access to the full design space: The bigger the design space the higher the likelihood to meet your goals in drug product formulation

Less time and material

By combining wet lab experiments with in-silico approaches, we find the right formulation with minimal use of drug substance in less time

Maximum flexibility

- Full formulation development, fast track and formulation optimization options available
- Choose which area of the design space is used

Formulation Technology Platform

- Library of up to 100 different regulatory established and employed excipients
- Rational development approach utilizing Bioinformatics, Biostatistics, State of the Art Design of Experiment (DoE) matrices

Advanced stabilization of a broad range of

- ✓ RNA
- ✓ LNP
- ✓ Exosomes
- ✓ Peptides
- ✓ Antibodies
- ✓ Proteins
- ✓ Viruses
- ✓ Vaccines
- ✓ Viral Vectors
- ✓ Enzymes

Specialized Services

- ✓ Molecular modelling
- ✓ Particle characterization
- ✓ Colloidal and conformational stability analysis
- ✓ Molecular purity and integrity
- ✓ Non-GMP drug product manufacturing
- ✓ Critical temperature analysis
- ✓ Lyo optimization
- ✓ In-silico long term stability prediction
- ✓ Formulation optimization analysis

Contact us and learn more:  bd@leukocare.com  www.leukocare.com



USA: Leukocare Inc., 27 Maple Street, Milford, MA 01757

Germany: Leukocare AG, Am Klopferspitz 19a, 82152 Martinsried/Munich

FUNDING OPPORTUNITIES IN INNOVATIVE PROJECTS FOR THE 2021–2027 PERIOD

Nikolina Letić

nikolina@lumennice.hr

Over the past couple of decades, the financing of research and development has been subject to significant changes. Public funding is not sufficient, traditional bank loans are generally difficult to obtain and yet there are EU funds that offer variety of possibilities for faculties and companies dealing with research and development to fund their innovative ideas at different level of development.

Croatian institutions and companies have access to variety of funding opportunities that can boost their research and improve their capacities in order to help them develop new processes, products or services, test innovative ideas and collaborate with other organizations. Funds that are allocated to Croatian institutions by the EU budget through *Multiannual Financial Framework 2021-2027* and through *Next Generation EU* are significant and are planned to be used to increase research and development activities in Croatian research institutions. Furthermore, research institutions and companies are entitled to apply for funding from *Horizon Europe* which is the EU's key funding programme for research and innovation for 2021 -2027 with a budget of €95.5 billion.

While average gross domestic spending on R&D in EU exceeded 2,32% in 2020, it is still rather low in Croatia - 1,25% in 2020. Thus, Croatian public institutions should build their capacities in order to use available funding opportunities.

REFERENCES

1. https://research-and-innovation.ec.europa.eu/funding_en

DISCOVERIES, CONTROVERSIES AND BEHEADINGS – HOW WATER STOPPED BEING AN ELEMENT

Vladimir Stilinović

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

vstilinovic@chem.pmf.hr

On November 12th 1783, at a public session of the French Royal Academy of Sciences, Antoine Laurent Lavoisier read out his memoir “*on the nature of water and on experiments which appear to prove that this substance is not strictly speaking an element but that it is susceptible of decomposition and recomposition*”.¹ In it he presented his discovery that water is in fact a chemical compound, comprising oxygen and hydrogen, and thus ended over two millennia of undisputed belief that water was an element. At least, that is the short story as usually told...

The exact nature of water was one of the central problems in the final decades of the phlogiston theory, and was tackled, with variable success, by many chemists of the day. The development of pneumatic chemistry during the 18th century has led many to observe connections between water (still considered an element in the light of the phlogiston theory) and the newly discovered (or at least identified and described) gases – mainly the dephlogisticated air (oxygen) and the inflammable air (hydrogen). This has led to several competing claimants to the priority of the discovery, and many of the controversies which have started in those days are still present amongst the historians of chemistry: has Lavoisier really made a profound discovery, or just changed the wording of Henry Cavendish; how much did Cavendish avail himself of the conclusions of James Watt; did Watt have any clearer idea than Joseph Priestly...²

Regardless of the questions of priority, the description of the composition of water by Lavoisier in terms of his *antiphlogistic* theory was to remove the greatest obstacle which stood in the path of its acceptance – the origin of the *inflammable air from metals* (hydrogen), described in detail by Cavendish in 1766, believed by many to be pure phlogiston – and allowed him to present a well-rounded alternative theory, virtually ending the phlogiston era. In this talk we shall attempt to reconstruct the unfolding of the events which have marked these, possibly the most dynamic, but certainly the most dramatic, decades in the history of chemistry.

REFERENCES

1. A. L. Lavoisier, *Observations sur la Physique* 1783, 23, 452.
2. T. E. Thorpe, *Nature* 1898, 57, 546.



A century of
dedication to health.

pliva.hr

STRUCTURAL REQUIREMENTS FOR *IN VITRO* AND *IN VIVO* ACTIVITY OF NOD2 AGONISTS

Žiga Jakopin

Faculty of Pharmacy, University of Ljubljana, Aškerčeva 7, SI-1000 Ljubljana, Slovenia

ziga.jakopin@ffa.uni-lj.si

The innate immune receptor NOD2 represents an important target for the development of structurally defined small molecule immunomodulators. Here, we present the investigation of the structure–activity relationships of several series of novel synthetic desmuramylpeptide NOD2 agonists. Extensive exploration of chemical space culminated in the discovery of NOD2 agonists with improved *in vitro* as well as *in vivo* adjuvant properties. Of note, an adamantane-moiety-featuring compound emerged as the most potent NOD2 agonist in its structural class to date. Further, pivotal structural elements that confer *in vivo* adjuvant activity in conjunction with a liposomal delivery system have also been identified. Our findings thus provide deeper insights into the structural requirements of desmuramylpeptides for NOD2-activation and highlight the potential use of NOD2 agonists as vaccine adjuvants or general immunostimulatory agents.

ACKNOWLEDGMENTS This work was supported by the Slovenian Research Agency grants (P1-0208, J3-9256 and P1-0420) and the Croatian Science Foundation (HrZZ) (Project No. 7387).

REFERENCES

- 1 Jakopin et al., *J Med Chem* 2012, 55, 6478.
- 2 Gobec et al., *Eur J Med Chem* 2016, 116, 1.
- 3 Gobec et al., *J Med Chem* 2018, 61, 2707.
- 4 Guzelj et al., *J Med Chem* 2021, 64, 7809.
- 5 Guzelj et al., *ACS Med Chem Lett* 2022, 13, 1270.

TAILORING THE LIQUID/LIQUID INTERFACE OF TRADITIONAL AND PICKERING GREEN EMULSIONS

Giuseppe Cappelletti

Dipartimento di Chimica, Università degli Studi di Milano, via Golgi 19, 20133 Milan, Italy

giuseppe.cappelletti@unimi.it

Emulsions are commonly employed in many different fields including pharmaceuticals, drug delivery, cosmetics, food industry, and so on, especially after the advancement of methods for preparing various kinds of emulsions.

Surfactants (or surface-active agents) are amphiphilic compounds widely used in almost every industrial field because of their ability to reduce surface and interfacial tensions, thus stabilizing dispersed systems. In recent years, due to their high worldwide consumption volumes, surfactant residues are becoming an environmental risk because they are continuously discharged into treated and untreated wastewaters, thus entering in several environmental compartments, i.e., surface and marine waters, sediments and sludge-amended soils etc.¹ Environmental concerns about effects of conventional tensides, mainly derived from petroleum precursors, have increased the market demand for more benign compounds. Particularly, sugar fatty acid esters (SFAEs)² belong to the non-ionic surfactant family, extensively adopted as emulsifiers in many market sectors, i.e. cosmetic, food and pharmaceutical industry, being low toxic and easy biodegradable. Thus, innovative syntheses of some glycoside esters will be proposed, and their surface properties/emulsifying capabilities will be studied in details.

Over the last decades, Pickering emulsions have attracted increasing attention due to their advantages compared to conventional surfactant-stabilized emulsions. Particularly, Pickering emulsions are characterized by long-term stability to coalescence, good mechanical properties, low toxicity and recyclable stabilizers³. In this context, Pickering emulsions represent promising platforms for applications in numerous fields, such as medicine, pharmaceuticals, cosmetics, food industry, and cultural heritage protection. In this work, highly stable and environmentally-friendly Pickering emulsion systems were prepared using food-grade vegetable oil and stabilized by the sole addition of pristine ZnO particles. The versatility of this approach was also proved changing the type of vegetable oil, the oxide content and morphology.

ACKNOWLEDGEMENTS This work was financially supported by Cariplo Foundation (Italy) (call: “Circular Economy for a sustainable future 2020”, project BioSurf, ID 2020-1094).

REFERENCES

1. M. Jackson, C. Eadsforth, D. Schowanek, T. Delfosse, A. Riddle, *Environ. Toxicol. Chem.* 2016, 35, 16.
2. S. Sangiorgio, E. Pargoletti, M. Rabuffetti, M. Robescu, R. Semproli, D. Ubiali, G. Cappelletti, G. Speranza, *Colloid Interface Sci. Commun.* 2022, 48, 100630.
3. C. Cionti, G. Vavassori, E. Pargoletti, D. Meroni, G. Cappelletti, *J. Colloid Interface Sci.* 2022, 628, 82.

CAS SCIFINDER DISCOVERY PLATFORM™ BETWEEN QUESTION AND ANSWER, THERE IS DISCOVERY

Accelerate breakthroughs and get discoveries to market faster using best-in-class scientific solutions throughout the innovation journey

Unmatched content

Specialized technology

Human expertise

Learn more at cas.org



ini INSTITUTE
FOR MEDICAL RESEARCH
AND OCCUPATIONAL HEALTH

Research and Educational Centre of Environmental
Health and Radiation Protection



ReC-IMI

75 YEARS OF EXPERIENCE IN
BASIC AND APPLIED RESEARCH



Environmental health assessment



Analytical and molecular toxicology



Dosimetry and radiation protection



Occupational health and psychology

RESEARCH & DEVELOPMENT ■ EDUCATION ■ SERVICES ■ PUBLISHING ■ SCIENCE POPULARIZATION



ALL BUT ONE: HOW NEGATIVE CATALYSIS SHAPED EVOLUTION OF THE ISOLEUCYL-TRNA SYNTHETASE'S EDITING DOMAIN

Igor Živković,* Aleksandra Maršavelski, Ita Gruić Sovulj

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* izivkovic.chem@pmf.hr

Aminoacyl-tRNA synthetases (AARS) activate amino acids and transfer them to cognate tRNAs. Some AARS cannot achieve the required specificity in the initial amino acid recognition at the synthetic site. Those AARS can incorrectly activate non-cognate amino acids and transfer them to tRNA. To correct the mistake and ensure faithful protein biosynthesis, these AARS evolved a separate editing domain aimed to hydrolyze erroneously aminoacylated tRNAs (post-transfer editing). To understand what shaped the selectivity of the editing site and how it is related to the initial amino acid selectivity at the synthetic site, we used *Escherichia coli* isoleucyl-tRNA synthetase (IleRS) as a model enzyme.^{1–3} We investigated the synthetic and editing reactions of IleRS using a wide range of amino acids, including non-proteinogenic and synthetic: Val, Nva, Leu, Thr, Met, Ser, Ala, Abu, Nle, F₂Abu and F₃Abu. Among these, only Val and Nva were poorly discriminated at the synthetic site (i.e. well activated and transferred to the tRNA) and thus can be considered as threat to the fidelity of translation. To our surprise, tRNAs misaminoacylated with all tested amino acids were rapidly hydrolyzed at the editing domain. Thus, it appears that amino acid's physicochemical features and how well it is rejected at the synthetic site does not pose a significant difference. Only the hydrolysis of cognate Ile-tRNA^{Ile} (misediting) was slow, suggesting that the need to keep the cognate product out of editing strongly shaped the specificity of the editing domain. Detailed kinetic analysis revealed that IleRS employs Thr246 and His333 for specific destabilization of Ile-tRNA^{Ile} hydrolysis – strategy also known as negative catalysis. Such design enabled IleRS to have broad substrate acceptance at the editing site whilst maintaining a high specificity towards preventing the futile post-transfer editing cycles. This was the first time such broad substrate specificity, paired with negative catalysis, was observed for an AARS, and as such, it marks a new moment in the understanding these vital and ancient enzymes.




Figure 1. Specificity and activity towards different substrates in the synthetic and editing sites of IleRS.

REFERENCES

1. Biluš M., et al, *Journal of Molecular Biology* 2019, 431, 1284.
2. Živković I., et al, *The FEBS Journal* 2020, 287, 800.
3. Živković I., et al, *Nucleic Acids Research* 2022, 50, 4029.

SECTION LECTURES

CYCLOPEPTIDES AS VERSATILE ANION SENSORS IN SOLUTION

Gordan Horvat

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

ghorvat@chem.pmf.hr

The design and synthesis of anion receptors is an ongoing hot topic in supramolecular chemistry. One especially interesting class of anion receptors are cyclopeptides. Their exceptional complexation properties are related to the proton-donating peptide groups, the flexibility of macrocyclic ring and to the variability of the macrocycle subunits. For these reasons, cyclopeptides stand as versatile receptors for halide and oxoanions in solution.

In this talk I will present our recent contributions to this field.¹⁻³ The aim of our research was to obtain as detailed as possible insight into the thermodynamic and structural characteristics of cyclopentapeptide complexes with halide and oxoanions in solution and to rationalize the relation between structure and reactivity of the investigated cyclopeptides. For the characterization of receptor-anion complexes we used various experimental methods, as well as molecular dynamics simulations. The solvent effect on equilibria of studied reactions was addressed as well.




Figure 1. Cyclopentaphenylalanine in complex with the chloride anion.

ACKNOWLEDGEMENTS This work was supported by the Croatian Science Foundation under project IP-2019-04-9560 (Macrosol) and European Regional Development Fund (project CluK, KK.01.1.1.02.0016).

REFERENCES

1. I. Petters, M. Modrušan, N. Vidović, I. Crnolatac, N. Cindro, I. Piantanida, G. Speranza, G. Horvat, V. Tomišić, *Molecules* **2022**, *27*, 3918.
2. G. Horvat, S. Tarana, N. Vidović, N. Cindro, G. Speranza, V. Tomišić, *J. Mol. Liq.* **2021**, *340*, 116848.
3. N. Vidović, G. Horvat, D. Riva, T. Rinkovec, N. Cindro, V. Tomišić, G. Speranza, *Org. Lett.* **2020**, *22*, 2129.

METALLOSUPRAMOLECULAR ASSEMBLIES – FROM SIMPLE COMPLEXES TO FUNCTIONAL NANOSTRUCTURES

Gracjan Kurpik,^{a,b,*} Anna Walczak,^{a,b} Grzegorz Markiewicz,^{a,b} Jack Harrowfield,^c Artur R. Stefankiewicz^{a,b}

^a Faculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland, ^b Center for Advanced Technology, Uniwersytetu Poznańskiego 10, 61-614 Poznań, Poland,

^c Institut de Science et d'Ingénierie Supramoléculaires, Université de Strasbourg, 8 allée Gaspard Monge, 67083 Strasbourg, France

* gracjan.kurpik@amu.edu.pl

Metal-ligand interactions are seen as one of the primary driving forces in the generation of simple coordination compounds, as well of metallosupramolecular architectures with a high degree of complexity and a precisely defined structure.¹ Taking into account the character of coordinate bond, i.e., high energy, but dynamic nature, coordination-driven self-assembly processes are characterized by the high controllability at the molecular level with respect to directionality, reversibility and then post-synthesized switchability.² In the presented research, a bifunctional ligand 4,4-dimethyl-1-(pyridin-4-yl)pentane-1,3-dione (HL) containing two distinct coordination sites, i.e., anionic β -diketonate and neutral pyridine has been used in the synthesis of Ag(I), Pd(II) and Pt(II) complexes that then have been applied as metalloligands for the construction of new heterometallic polymeric materials. The ambidentate nature of HL enables switching between different modes of coordination within mononuclear complexes or their conversion into polymeric species in a fully controllable way. The coordination-driven processes can be triggered by various stimuli, i.e., a metal salt addition or acid-base equilibria, and presents an efficient strategy for the generation of metallosupramolecular materials. As a consequence of self-assembly, new heterometallic coordination aggregates have been synthesized and fully characterized via different analytical techniques in solution as well as in the solid state.

AMBIDENTATE LIGAND HL




Figure 1. The supramolecular transformations within coordination-driven assemblies based on Ag(I), Pd(II) and Pt(II) ions and the pyridyl- β -diketonate ligand HL.

ACKNOWLEDGEMENTS The financial support was provided by the National Science Centre (ARS, grant SONATA BIS 2018/30/E/ST5/00032).

REFERENCES

1. E. C. Constable, C. E. Housecroft, *Chem. Soc. Rev.* **2013**, *42*, 1429.
2. W. Wang, Y.-X. Wang, H.-B. Yang, *Chem. Soc. Rev.* **2016**, *45*, 2656.

Molekulska spektroskopija

UV/Vis spektrofotometrija

- mali, prijenosni spektrofotometri
- laboratorijski uređaji Lambda serije



Lambda 265



Lambda 365



Lambda 465



Lambda XLS

FTIR spektroskopija



Spectrum Two



Frontier



Spotlight 150/200i/400



Kromatografija

- plinska i tekućinska kromatografija



Clarus GC



Flexar



LC 300



GC 2400 Platform

Anorganske analize

- atomska spektrometrija
- ICP-OES
- ICP-MS



AAS PinAAcle serija



ICP-OES Avio serija

ICP-MS

NextION serija



Fluorimetrija



FL6500

FL8500

Ostala laboratorijska oprema



- uređaji za mikrovalnu digestiju
- uređaji za pročišćavanje kiselina
- niskotlačni / visokotlačni reaktori



SPEX SamplePrep

SPEX CertiPrep

Organic & Inorganic Carbon Dioxide Materials

- homogenizatori, kriogeni mlinovi, kuglični mlinovi
- standardi CRM (organski, anorganski, „custom made“)

KONTAKT:

Gabriela Lihtar, mag.chem.

E-mail: gabriela.lihtar@jasika.hr

Marijana Jurić, mag.chem.

E-mail: marijana.mihoci@jasika.hr



We create comprehensive solutions for reliable and credible analyses.



- Laboratory equipment
- Application support
- Service




FROM SINGLE MOLECULE TO RESPONSIVE AGGREGATE: MULTIRESPONSIVE PHOTOSWITCH IN NONPOLAR SOLVENTS

Miroslava Čonková,^{a,b,*} Łukasz Majchrzycki,^b Artur R. Stefankiewicz^{a,b}

^aFaculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland, ^bCenter for Advanced Technology, Adam Mickiewicz University, Uniwersytetu Poznańskiego 10, 61-614 Poznań, Poland.

* miroslava.conkova@amu.edu.pl

Multi-responsive, also known as smart, materials are currently one of the most appealing part of materials science.¹ Smart polymers are able to change macroscopically their structure, but also these transitions are reversible and the system return to the original state after the stimulus is removed. Since the 1990s when smart materials first appeared in the literature,² different reversible transformations were observed. Our unique design combines the simplicity of a monomer (low molecular weight with well-defined primary structure) with the complex properties of self-assembled architecture. The special focus is on the synergism of several different responses of the architecture, which are encoded in different moieties of the monomer. Herein we present simple yet robust tetra substituted azobenzene molecule, bearing C_4 symmetry. When dissolved in a polar solvent, the molecule behaves as a monomer and is fully molecularly dissolved. On the other hand, when in a nonpolar solvent, supramolecular aggregation occurs, forming large aggregates via hydrogen bonds and π - π stacking. After heat or light are employed we are able to control the size of the aggregate, however, when a different type of stimulus is present (acid, acetonitrile or combination of heat and solvent) even full disassembly of the aggregate is achieved. The aggregate and its behavior after applied stimuli was monitored in solution by several spectroscopic methods and on the surface by AFM.



ACKNOWLEDGEMENTS The project was funded by the National Science Centre of Poland grants SONATA BIS 2018/30/E/ST5/00032. The conference attendance was funded by grant "Środowiskowe interdyscyplinarne studia doktoranckie w zakresie nanotechnologii" No. POWR.03.02.00-00-I032/16.

REFERENCES

1. L. Peponi, M. P. Arrieta, A. Mujica-Garcia, D. López, in *Modification of Polymer Properties* (Ed.: C. F. Jasso-Gastinel, Kenny J. M.), Elsevier Inc, Oxford, United Kingdom, **2017**, pp. 131-154.
2. C. Fouquey, J.-M. Lehn, A.-M. Levelut, *Adv. Mater.* **1990**, *2*, 254.

SOLUTIONS FOR POLLUTANTS SENSING – SURFACTANTS, ESTROGEN AND GLYPHOSATE SENSORS

Nikola Sakač^{a,*} Marija Jozanović,^b Dubravka Madunić-Čačić,^c Bartolomeo Della Ventura,^d
Raffaele Velotta^d

^a Faculty of Geotechnical Engineering, University of Zagreb, 42000 Varaždin, Croatia,

^b Department of Chemistry, University of Osijek, 31000 Osijek, Croatia, ^c Saponia Chemical,
Pharmaceutical and Foodstuff Industry, Inc., 31000 Osijek, Croatia,

^d Department of Physics “E. Pancini” – Università Di Napoli Federico II, 80126 Naples, Italy

* nsakac@gfv.unizg.hr

Chemical sensors and biosensors are analytical devices that perform a specific or selective recognition (recognition element) of target analyte with transducer that converts the recognition into a readable signal. They have a wide range of usage, from environmental, industrial and biomedical, up to military. Global chemical sensor market in 2020 was USD 21.39 billion with estimated growth 7.51% from 2020 to 2026¹, and USD 24.9 billion in 2021 for biosensor market with estimated growth 8% from 2022 to 2030². The main focus in development of chemical (bio)sensors is to be low-cost, easy to use, sensitive and selective/specific, robust, small and portable. The lecture will cover the research results of the development and application of electrochemical potentiometric anionic and cationic surfactant sensors based on the PVC-liquid membrane type membranes^{3,4}, and nanoparticle based localized surface plasmon resonance (LSPR) optical sensors for estrogen hormone 17 β -estradiol⁵ and LSPR optical immunosensor with magnetic micromixing extraction for highly sensitive detection of a broad-spectrum systemic herbicide glyphosate⁶. The research was recently conducted at the Laboratory for environmental engineering at the Faculty of geotechnical engineering, University of Zagreb (Croatia) in cooperation with partners from home and international research institutions and industry.

Keywords: Surfactant, glyphosate, estradiol, chemical sensor, biosensor

REFERENCES

1. mordorintelligence.com, visited: 1 October 2022
2. grandviewresearch.com, visited: 1 October 2022
3. Sakač, N., Marković, D., Šarkanj, B., Madunić-Čačić, D., Hajdek, K., Smoljan, B., Jozanović, M. *Molecules* 2021, 26, 1420.
4. Sakač, N., Madunić-Čačić, D., Marković, D., Hok, L., Vianello, R., D., Šarkanj, B., Đurin, B., Hajdek, K., Smoljan, B., Milardović, S., Matasović, B., Jozanović, M. *Molecules* 2021, 14, 3627.
5. Minopoli, A., Sakač, N., Lenyk, B., Campanile, R., Mayer, D., Offenhäusser, A., Velotta, R., Della Ventura, B. *Sensors and Actuators B: Chemical* 2020 308, 127699.
6. Campanile, R., Elia, C., Minopoli, A., Ud Din Babar, Z., Di Girolamo, R., Morone, A., Sakač, N., Velotta, R., Della Ventura, B., Iannotti, V. *Talanta* 2023 253, 123937.

Bringing the Power of AI to Analytical Laboratories

NEXERA UHPLC



A New Benchmark of **Intelligence**
Maximizing Reliability, Minimizing Down Time

Shimadzu d.o.o. - Zavrtnica 17 - 10000 Zagreb - tel. +385 1 6185 777

shimadzu@shimadzu.hr - www.shimadzu.hr



Znanstvena i laboratorijska rješenja na polju:

- Kemikalija i reagensa
- Mikrobiologije
- Molekularne biologije, proteina i stanične kulture
- Sirovina i bioprocasa
- Laboratorijske plastike

Kontakt

Tel: 01 4800 111

Mail: kemikalije@medic.hr

Web: www.medic.hr

Naši brandovi:

MERCK

MILLIPORE SIGMA



greiner
BIO-ONE

Luminex®

lipomed
SERVICES TO HEALTH

TRIARYLBORANES: NOVEL TYPE OF DYE FOR BIOMACROMOLECULES


Ivo Piantanida,^{a,*} Željka Ban,^a Marta Koščak,^a Todd B. Marder^b

^a Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička 54, Zagreb, Croatia,

^b Institut für Anorganische Chemie, and Institute for Sustainable Chemistry & Catalysis with Boron, Julius-Maximilians-Universität Würzburg, Würzburg, Germany

* pianta@irb.hr

Low molecular weight fluorescent probes are frequently used tools in most molecular biology or biochemistry experiments today.¹ Due to problems related to covalent attachment of dye to biomolecule, non-covalently binding molecular probes that readily recognize and distinguish between different types of biomolecules are of huge interest.² However, the small size of a molecule presents a significant challenge in the design of selective probes, particularly in well-explored fluorophore families. Thus, development of a novel structural motif represents a valuable contribution. Here we report the development of a novel family of dyes based on the triarylborane system (Scheme 1.), which showed remarkable features in the interaction with biomacromolecules (DNA, RNA, proteins).



Scheme 1. Some applications of triarylborane dyes: LEFT: fluorescent response differing between DNA and protein,³ or RIGHT: dual sensing of DNA or protein by fluorescence and SERS.⁴

Namely systematic structural variation (mostly linkers between two triarylborane units) revealed high tunability of a system in respect to emission response and Raman (SERS) response, promising photo-bioactivity or intriguing enzyme-staining properties.

ACKNOWLEDGEMENTS The financial support of the Croatian Science Foundation project IP-2018-01-5475, the DAAD, and the Julius-Maximilians-Universität Würzburg are gratefully acknowledged.

REFERENCES

1. R. B. Silverman, in *The Organic Chemistry of Drug Design and Drug Action*, Elsevier Academic Press, Amsterdam, 2nd edn., 2004.
2. M.-A. Mycek, B. W. Pogue, in *Handbook of Biomedical Fluorescence*, CRC Press, 2003.
3. Z. Ban, S. Griesbeck, S. Tomic, J. Nitsch, T. B. Marder, I. Piantanida, *Chem. Eur. J.* 2020, 26, 2195.
4. M. Ferger, Z. Ban, I. Krošl, S. Tomić, L. Dietrich, S. Lorenzen, F. Rauch, D. Sieh, A. Friedrich, S. Griesbeck, A. Kendel, S. Miljanić, I. Piantanida, T. B. Marder, *Chem. Eur. J.* 2021, 27, 5142.

JGL'S EXPERIENCE IN DEMONSTRATING EXTENDED PHARMACEUTICAL EQUIVALENCE IN DRUG PRODUCT SOLUTIONS

Maša Safundžić Kučuk,^{a,*} Leo Štefan,^a Zdravka Knežević^b

^a Research and Development, JGL d.d., Svilno 20, Rijeka, Croatia, ^b Scientific Operations, JGL d.d., Svilno 20, Rijeka, Croatia

* masa.safundzic@jglpharma.com

During development of a pharmaceutical generic product, demonstration of its qualitative (Q1), quantitative (Q2), and microstructural (Q3) equivalence to the reference medicinal product (RMP) is the cornerstone of the R&D activities. Apart from the legal requirement to demonstrate quality, safety, and efficacy similarity with the RMP, generic companies invest considerable effort to demonstrate Q1/Q2/Q3 equivalence in order to be exempted from the costly and lengthy clinical trials, or to limit their extent.¹ First step in the demonstration of the equivalence, is to fully characterize the RMP. Full composition of the RMP is rarely fully known, and as guidelines allow up to $\pm 5\%$ difference in composition for a generic product from the RMP,² to consider a Q1 and Q2 equivalence demonstrated (some exceptions are allowed). To achieve this, a developer must combine extensive literature research and diverse reverse engineering activities to determine exact composition of the RMP. If a company decides to develop a product that deviates from the Q1/Q2 equivalence – for example, by excluding preservatives, or to bypass a valid patent, status of a generic product still can be achieved with demonstration of the Q3 equivalence.³ In demonstration of the latter, various techniques are used for the purpose of characterization of the critical quality attributes (CQA) and their similarity with RMP. For each project, a case-by-case basis approach is needed, and to obtain reliable results and conclusions, in-depth knowledge of the pharmaceutical technology, analytical, physical, and organic chemistry is essential. For the purpose of accessing the specialized knowledge, or to accelerate development process, Companies often outsource their development studies. In some cases, Academia seems like an ideal partner, but some requirements need to be met.

ACKNOWLEDGEMENTS Vesna Saršon, Silvia Kamhi Saršon

REFERENCES

1. European Medicines Agency, *Note for Guidance on the Clinical Requirements for Locally Applied, Locally Acting Products Containing Known Constituents*, 1995, CPMP/EWP/239/95 final.
2. European Medicines Agency, *Draft Guideline on quality and equivalence of topical products*, 2018, CHMP/QWP/708282/2018.
3. V. P. Shah, A. Yacobi, F. S. Rădulescu, D. S. Miron, M. E. Lane, *International Journal of Pharmaceutics*, 2015, 491, 21.

Već više od 30 godina tvrtka **AnAs d.o.o.** snabdijeva analitičkom opremom i pripadajućim priborom brojne znanstveno – istraživačke institucije i industrijske pogone u Republici Hrvatskoj.

Široka paleta proizvoda koje nudimo, u kombinaciji s našim znanjem i iskustvom stečenim radom u laboratoriju, omogućuje nam da ponudimo najbolje rješenje za sve vaše analize.

Jedno od naših temeljnih načela jest etikom vođeno poslovanje, jer se tako zadovoljavaju ključne vrijednosti: vjerodostojnost, pouzdanost, kompetentnost i čestitost.

Na zadovoljstvo naših korisnika kontinuirano unapređujemo i prilagođavamo poslovanje njihovim potrebama, ostvarujući time svoj cilj.

analytikjena
An Endress+Hauser Company

CEM

scion
INSTRUMENTS

BRUKER

TELEDYNE ISCO
Everywhere you look™

Rigaku

RESTEK

VICI

VICI DBS

CPA chem

INORGANIC VENTURES
Refine your results. Redefine your industry.

Starna scientific

hta
sampling for science

Organomation

Specac

GL Sciences

Svojim korisnicima nudimo samo proizvode najviše kvalitete, stoga nam se obratite s povjerenjem!

bia

ZANESLJIV
LABORATORIJSKI
PARTNER

TRUSTED LAB
PARTNER

BIA d.o.o. (Biotechnology, Instrumentation, Analysis)

Već više od 30 godina distribuiramo vodeće svjetske proizvođače kao što su:

- Advion Interchim Scientific (flash i preparativna kromatografija)
- Distek (dissolution oprema, BIONe Single-Use Bioreactor (SUB))
- Knauer (analitička, preparativna i SMB kromatografija, FPLC, LNP)
- Infors HT (inkubatorski shakeri, bioreaktori)
- Radleys (oprema za kemijsku sintezu)
- SP Genevac

i mnogo više..... potraži na www.bia.si;
sales@bia.si; 00385916011720

CAUSAL ANALYSIS OF MOLECULAR DESCRIPTORS OF EUTECTIC SOLUTIONS OF CHOLINE CHLORIDE SYSTEMS APPLICABLE IN PHARMA TECHNOLOGIES

Želimir Kurtanjek

Faculty of Food Technology and Biotechnology, University of Zagreb, Zagreb, Croatia

zelimir.kurtanjek@gmail.com

Applied is the structural causality model (SCM) analysis of molecular descriptors and fingerprints for predicting the eutectic temperature of choline chloride systems aimed for application in pharmaceutical technologies. The SCM model predictions are compared to predictions by the ideal mixture model and COSMO-RS (Conductor like Screening Model for Realistic Solvents). The model is developed by use of a dataset of 34 molecules including common pharmaceutical compounds (ibuprofen, ketoprofen, and paracetamol). The molecular descriptors and fingerprints generated by PaDEL software currently calculate 1875 descriptors (1444 1D, 2D descriptors, and 431 3D descriptors) and 12 types of fingerprints (a total of 16092 bits). The descriptors and fingerprints are calculated using The Chemistry Development Kit with additional descriptors and fingerprints such as atom-type electron topological state descriptors.¹⁻⁴ The large data sets are analyzed by the suit of numerical and statistical packages available in R software.⁵ The model of the directed acyclic graph (DAG) inferred is by Hamilton-Schmidt independence criteria (HSIC) The causal functionalities between eutectic temperature and molecular descriptors are determined by deconfounding by the d-separation criteria.⁶ The accuracy of SCM-DM is compared with COSMO-RS and is presented in Fig. 1. Mean absolute errors are 7.5 °C and 9.57 °C for SCM-DM and COSMO-RS respectively.⁷ The causal SCM model has important advantages for application in the scan evaluation of large molecule datasets in the development of new green sustainable technologies.




Figure 1. Comparison of eutectic temperature predictions by the causal structural SCM-MD and COSMO-RS real solution models.

REFERENCES

1. R. Todeschini, V. Consonni, "Handbook of Molecular Descriptors", **2000**, Wiley-VCH, USA.
2. C.W. Yap, *J. Comput. Chem.* **2011**, 32, 1466.
3. PubChem System <http://pubchem.ncbi.nlm.nih.gov>
4. D.O. Abranches et al., *Fluid Phase Equilibria*, **2019**, 497, 71.
5. R version 4.0.3, **2020**, <https://www.r-project.org>
6. J. Pearl, Causality, 2nd edition, **2021**, Cambridge University Press, UK.
7. Z. Kurtanjek, **2022**, *Chem. Biochem. Eng. Q.* (submitted).

SPECTROPHOTOMETRIC KINETIC AND THERMODYNAMIC INVESTIGATION FOR THE OXIDATION OF ANILINES BY HEXACHLOROIRIDATE(IV) IN SOLUTION MEDIA: A COMBINED EXPERIMENTAL AND COMPUTATIONAL STUDY

Riya Sailani

Department of Chemistry, Centre of Advanced Study, University of Rajasthan, Jaipur,
Rajasthan-302004 India

I.p_riya@yahoo.co.in

The oxidation kinetics of ortho and para substituted anilines with hexachloroiridate (IV) in aqueous perchloric acid have been performed. The reaction is indicated first order dependence with respect to both reactants. Reaction rates accelerated with the introduction of electron-releasing groups and retarded with electron-withdrawing groups which are elucidated by Hammett's theory of linear free energy relationship. Hammett's reaction constant (ρ) has negative ($\rho < 0$) value. Furthermore, an increase in temperature increased the reaction constant (ρ). Various thermodynamic parameters have been reported and discussed the validity of isokinetic relationship. Isokinetic temperature (β) from Vonthoff's plot (317 K), Compensation plot (317.4 K) and Arrhenius plot (317 K), is observed. Observed β value from Exner's plot (439.32 K) is above the experimental temperature range (303–323 K), indicating that the enthalpy factors are probably more important in controlling the reaction. The oxidation product is corresponding 2-keto azoxy benzenes. Based on the kinetic results, a suitable mechanism has been proposed.

To further support our proposed mechanism, density functional theory (DFT) computations were done at M06/6-311*G showing that activation energy barriers predict the same reactivity trend as shown by the kinetics experiments.

ACKNOWLEDGEMENTS This work was supported in part by the University Grant Commission, New Delhi, India through start-up grant (BSR) for financial support. Acknowledgment is also made to Head of Department of Chemistry, UOR, Jaipur, Rajasthan, India. Author wants to dedicate this work to (late) Prof. P. D. Sharma, ex-Head of Department of Chemistry, UOR, Jaipur, Rajasthan, India.

REFERENCES

1. R. Sailani, *ChemistrySelect*, **2021**, *6*, 8472.
2. R. Sailani, D. Pareek, A. Meena, K. Jangid, C. L. Khandelwal, *Ind. J. Chem.* **2016**, *55*, 1074.
3. O. Exner, *Nature* **1964**, *201*, 488.
4. L. Liu, Q-X. Guo, *Chem. Rev.* **2001**, *101*, 673.
5. S. S. Mansoor, S. S. Shafi, S. Z. Ahmed, *Arab. J. Chem.* **2017**, *10*, S1129.
6. N. E. Guesmi, T. Boubaker, R. Goumont, *Int. J. Chem. Kinet.* **2010**, *42*, 203.
7. R. Ibarz, A. Garvin, A. Ibarz, *Food Res. Int.* **2017**, *99*, 348.



Unsere Chemikalien-Marken: Qualität aus eigener Produktion



ML₂ METAL COMPLEXES WITH BPA AND IMDA LIGANDS – IN SOLUTION AND SOLID-STATE

Aleks Logožar,^{a,*} Natalija Pantalon Juraj,^b Berislav Perić,^b Robert Vianello,^b Srećko I. Kirin^b

^a Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a 10000 Zagreb

^b Division of Materials Chemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb

* alogozar@chem.phf.hr

Bis-tridentate metal complexes of flexible ligands with [M(A-X-A)₂] structure can form three different geometric isomers: *mer*, *trans-fac* and *cis-fac*.¹ Many factors influence the stability of a certain isomer. Important factors to consider are electronic and steric properties of the coordinating ligand, type of central metal ion and counter-ion present, possibilities of forming non-covalent interactions and others. Understanding of the isomer preferences in hereby described systems would give us a head start in the development of new coordination compounds with desired form and properties.^{2,3}

Derivatives of tridentate ligands bis(pyridine-2-ylmethyl)amine (bpa) and 2,2'-iminodiacetamide (imda) were prepared in this work, as well as their corresponding complexes with different Zn(II), Cu(II), Ni(II) and Co(II) salts (Figure 1). Oxygen and sulfur, as well as -NH-, -NPh-, -PH- and -PPh- functional groups were chosen to occupy a central position in the ligand structure. The stoichiometry and the stereochemistry of these complexes, together with possible conformations of their substituents, were studied using the DTF computational approach to determine their relative stability in solution. The systems in solution were studied by NMR and UV-Vis spectroscopy, while the solid-state structures of prepared complexes, characterized by single-crystal X-ray diffraction, gave us an insight of the isomerism in the solid-state. Experimental and computational results were discussed and preferences of some central donor atoms to form certain isomers of ML₂ complexes were shown. One of our goals was to determine which factors lead to formation of *cis-fac* isomers, which could, in the case of nitrogen or phosphorous as central atoms, be used in the development of new type of selective catalysts.




Figure 1. ML₂ complexes of flexible tridentate bpa and imda ligands.

ACKNOWLEDGEMENTS This project was partially supported by CAT Pharma (KK.01.1.1.04.0013).

REFERENCES

1. N. Pantalon Juraj, S. I. Kirin, *Coord. Chem. Rev.* **2021**, *445*, 214051.
2. N. Pantalon Juraj, G. I. Miletić, B. Perić, Z. Popović, N. Smrečki, R. Vianello, S. I. Kirin, *Inorg. Chem.* **2019**, *58*, 16445.
3. Đ. Škalamera, E. Sanders, R. Vianello, A. Maršavelski, A. Pevec, I. Turel, S. I. Kirin, *Dalton Trans.* **2016**, *45*, 2845.


HALOGENIDE ANIONS AS HYDROGEN AND HALOGEN BOND ACCEPTORS IN SIMPLE COCRYSTALS WITH A NEUTRAL HALOGEN BOND DONOR

Marin Liović,* Nikola Bedeković, Vladimir Stilinović, Dominik Cinčić

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* mliovic@chem.pmf.hr

In comparison to the other Lewis bases, halogenide anions stand out with having no covalently bound atoms which would hinder the approach of one or multiple Lewis acids. That being said, halogenides have been investigated in solid state as acceptors of either hydrogen or halogen bonds,^{1,2} but scarcely researched as simultaneous acceptors of both interactions. One of the simplest systems used for such an investigation are halogenopyridinium halogenides,³ as well as their cocrystals with neutral halogen bond donors.⁴ In order to examine halogenides in that role, we prepared cocrystals of 3,5-lutidinium and 2,4,6-collidinium halogenides (X = F, Cl, Br, I) with 1,4-diiodotetrafluorobenzene (**14tfib**) *via* solution synthesis. The molecular and crystal structures of seven cocrystals were determined from the data obtained by single crystal X-ray diffraction experiments. In all the prepared solids, along with N–H···X[−] hydrogen bonds formed between cations and halogenides, every halogenide and **14tfib** alternate forming 1D chains *via* C–I···X[−] halogen bonds. Comparing interaction lengths relative to the sum of van der Waals and ionic radii of the interacting atoms within both cation series, hydrogen bonds are relatively shorter for smaller halogenides while halogen bonds vary slightly in relative length and do not follow any trend. Despite the differences in cations and halogenides, halogen bonded motifs are preserved in all the cocrystals.



Schematic representation of cocrystal synthesis and the resulting motifs.

REFERENCES

1. G. Gilli and P. Gilli, *The Nature of the Hydrogen Bond*, Oxford University Press, Oxford, 2009.
2. G. Cavallo, P. Metrangolo, R. Milani, T. Pilati, A. Priimagi, G. Resnati, G. Terraneo, *Chem. Rev.* 2016, **116**, 2478.
3. L. Fotović, V. Stilinović, *CrystEngComm* 2020, **22**, 4039.
4. L. Posavec, V. Nemeč, V. Stilinović, D. Cinčić, *Cryst. Growth. Des.* 2021, **21**, 6044.



BIOCentre

incubation centre
for biosciences

The place where Science and Entrepreneurship successfully meet

Services:

- Microbiology
- Downstream Processing
- Bioanalytics – Chemistry
- Bioanalytics – Proteomics
- Cell Biology
- Workshops and seminars



BICRO BIOCENTRE, Ltd. • FOR TRANSFER AND COMMERCIALIZATION OF BIOTECHNOLOGY
 Borongajska cesta 83 H, 10 000 Zagreb, Croatia • t: +385 1 6458 640 • e: info@biocentre.hr • biocentre.hr



**NOVI
RAMAN**

METTLER TOLEDO



**NOVI
FTIR**

Moćna analiza procesa. Pojednostavljeno.

Snažna, robusna i jednostavna za korištenje
Spektroskopija za brže donošenje sigurnih odluka

Kako biste saznali više o našim novim FTIR i Raman tehnologijama, zatražite demonstraciju ili ponudu:

- ▶ www.mt.com/ReactIR
- ▶ www.mt.com/ReactRaman



APPLICATIONS OF POLYELECTROLYTE MULTILAYERS FOR THE REDUCTION OF BACTERIAL ADHESION

Davor Kovačević^{a,*} Klemen Bohinc^b

^a Division of Physical Chemistry, Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia, ^b Faculty of Health Sciences, University of Ljubljana, Zdravstvena pot 5, 1000 Ljubljana, Slovenia

* davor.kovacevic@chem.pmf.hr

Polyelectrolyte multilayers (PEMs) are very well known surface coatings which could be prepared by alternating deposition of positively and negatively charged polyelectrolytes (polycations and polyanions) on a solid surface. PEM build-up strongly depends on applied experimental conditions such as ionic strength, supporting electrolyte type, pH and concentration. In recent years we have developed new PEM strategies which could be valuable for designing soft nano materials whose properties can be finely tuned according to the requirements of specific applications, especially in the field of biomedicine. The emphasis in our studies was in the application of PEMs for the prevention of bacterial adhesion to various surfaces. We investigated the influence of polyelectrolyte multilayer properties on bacterial adhesion capacity,¹ as well as the bacterial adhesion capacity of protein-terminating polyelectrolyte multilayers.² We also applied this PEM strategy to study bacterial adhesion capacity of uropathogenic *Escherichia coli* to polyelectrolyte multilayer coated urinary catheter surface.³ In this case it was shown that on non-treated PVC surfaces, biofilm was formed which was not the case for polyelectrolyte multilayer coated surfaces.

ACKNOWLEDGEMENTS This research was supported by the Croatian Science Foundation under the bilateral Slovenian-Croatian APPLPEMS project (IPS-2020-01-6126).

REFERENCES

1. D. Kovačević, R. Pratnekar, K. Godič Torkar, J. Salopek, G. Dražić, A. Abram, K. Bohinc, *Polymers* **2016**, *8*, 345.
2. K. Bohinc, J. Bajuk, J. Jukić, A. Abram, M. Oder, K. Godič Torkar, P. Raspor, D. Kovačević, *Int. J. Adhes. Adhes.* **2020**, *103*, 102687.
3. K. Bohinc, L. Kukić, R. Štukelj, AM. Zore, A. Abram, T. Klačić, D. Kovačević, *Coatings* **2021**, *11*, 630.

COMPARISON OF ALBUMIN AND CHITOSAN EFFECTS ON CALCIUM PHOSPHATE FORMATION ON TITANATE NANOMATERIALS

Ina Erceg,* Maja Dutour Sikirić

Laboratory for Biocolloids and Surface Chemistry, Division of Physical Chemistry,
Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia

* ierceg@irb.hr

The constant need for advanced materials for bone regeneration motivates the development of calcium phosphate (CaPs) composites with different types of materials.¹ The most frequently used materials are biomacromolecules for enhancing biological response and nanomaterials for improving mechanical properties. A possible method for preparing such composites is precipitation at low temperatures,^{2,3} a simple and inexpensive method that allows successful incorporation of components with low stability. Recently, the composites of CaPs, biomacromolecules and nanoparticles have emerged as promising new materials for bone regeneration. To successfully prepare such multicomponent materials by precipitation, a study of the influence of individual components on CaPs formation is required.

In this study, the effects of bovine serum albumin (BSA) and chitosan (Chi) on the precipitation of CaPs on titanate nanomaterials (TiNMs) of different morphologies (nanoparticles, nanoplates, nanotubes, and nanowires) were compared. Both biomacromolecules inhibited the transformation of amorphous to the crystalline phase, regardless of the presence of TiNMs which promote the transformation. Neither biomacromolecule influenced the composition of the formed CaP, as calcium deficient hydroxyapatite (CaDHA) was formed in all cases. However, in the presence of both BSA and Chi, the crystalline size and CaDHA morphology were influenced. As well, they inhibited the formation of CaDHA on the surface of the nanowires.

The obtained results may contribute to the development of precipitation methods for the preparation of multicomponent composites.

ACKNOWLEDGEMENTS Financial support from Croatian Science Foundation, Grant HRZZ- IP-2018-01-1493 is greatly acknowledged.

REFERENCES

1. S. V. Dorozhkin, *J. Mater. Chem. B* **2019**, *7*, 7471.
2. K. Duan, R. Wang, *J. Mater. Chem.* **2006**, *16*, 2309.
3. I. Erceg, A. Selmani, A. Gajović, I. Panžić, D. Iveković, F. Faraguna, S. Šegota, M. Ćurlin, V. Strasser, J. Kontrec, D. Kralj, N. Maltar Strmečki, M. Dutour Sikirić, *Colloids and Surfaces A: Physicochemical and Engineering Aspects* **2020**, *593*, 124615.

PROTECTIVE ABILITY OF *ARTEMISIA ANNUA* L. AGAINST MICROBIOLOGICALLY INFLUENCED CORROSION OF ALUMINIUM ALLOY 5083 CAUSED BY *PSEUDOMONAS AERUGINOSA* IN ARTIFICIAL SEAWATER

Gloria Zlatić,^a Ivana Martinović,^a Zora Pilić,^a Stipe Čelan^b

^a Department of Chemistry, Faculty of Science and Education, University of Mostar, Matice hrvatske bb, 88000 Mostar, ^b Institute for Food and Veterinary of Herceg-Bosnian Canton, Obrtnička 2, 80101 Livno

* gloria.zlatic@fpmoz.sum.ba

Aluminium-magnesium alloys, 5000 series, are used in many marine applications because of their versatile benefits of strength, weldability, and high corrosion resistance. The mechanisms of corrosion processes in marine environments are quite complex and largely dependent on pH, concentration of dissolved gases, temperature, salinity and the presence of microorganisms. Certain microorganisms are able to colonize the metal surface and indirectly inhibit or accelerate corrosion processes. Microbiologically influenced corrosion (MIC) of aluminium alloy 5083 (ALA) caused by the bacterium *Pseudomonas aeruginosa* (PA) in artificial seawater (ASW) was investigated by electrochemical, surface and spectroscopic techniques without and with the addition of *Artemisia annua* L. extract (AAE). Electrochemical tests proved good corrosion resistance of ALA in ASW, where corrosion current densities, j_{corr} determined from Tafel plots decreased with the increasing immersion time, which was confirmed by electrochemical impedance spectra (EIS) recorded at open circuit potential (OCP) where an increase in the resistance of the film was observed. In a medium inoculated with PA, OCP of ALA shifted to more negative values, while obtained j_{corr} values increased with the increasing immersion time which indicated that PA accelerated corrosion of ALA in ASW. Further obtained results showed that the plant extract prevents the MIC of ALA in ASW by reducing both anodic and cathodic current densities with an inhibition efficiency of 80 %. After 14 days of incubation, the addition of plant extract to inoculated medium at a concentration of 1.0 g L⁻¹, led to an increase in resistance of the surface film, which confirmed inhibitory effect of AAE on MIC of ALA. 3D profiles of treated ALA surface, obtained by Optical Profilometer (OP) revealed that pitting of ALA occurred during 21 days incubation in ASW inoculated with PA, while the addition of AAE led to a decrease in film thickness and roughness. The concentration of aluminium ions in the solution, measured by inductively coupled plasma - optical emission spectrometry (ICP-OES) after short immersion tests of treated coupons, was also in accordance with the electrochemical results, whereby the measured concentration in the inoculated samples was 77.6 µg L⁻¹ cm⁻², while after the addition of AAE, the concentration was 43.5 µg L⁻¹ cm⁻².

ACKNOWLEDGEMENTS FBiH Ministry of Education and Science, FBiH Agro-Mediterranean Institute, CEEPUS

INNOVATIVE APPROACHES IN TEACHING CHEMISTRY

Daria Stejskal

OŠ Frana Galovića, Zagreb, Croatia

* dariastejskal@gmail.com

The physiology of learning and memory of today's children is different. A higher level of activity, speed, and stimulus is needed. There is an ever-widening gap between students' needs and established teaching methods, which causes dissatisfaction on both sides. Students are unmotivated, they do not see the point of learning, and they give up further schooling, especially in the field of science. On the other hand, regardless of major advances achieved in the approach to teaching and in the light of new insights in the field of neuroscience and neurodidactics, teachers ought to adapt to new generations and change learning strategies by combining formal and informal methods. According to findings from neuroscience, learning should be meaningful, connected with actual situations (contextual), directed towards learning by solving problems, research, cooperative learning, and play-based learning. The student is thus placed at the centre of learning, while the teacher acts as a moderator who directs and encourages the process itself.¹ Using the stated learning strategies, we do not separate emotions from cognition, which neuroscientists find to be decisive in the learning process.

Information „coloured“ by emotion has a greater meaning, i.e. according to the broaden-and-build theory, positive emotions build and broaden the repertoires of thoughts and actions. In that context, students learn by creating and constructing their own insights (knowledge) based on interaction with the environment (physical and social) and by imparting their own meaning, understanding, and interpretation to the environment. Active learning strategies; practical work, research, and games affect the plasticity of the brain by forming new neural connections.²

Taking into account the mentioned findings and needs, both of students and teachers, at Fran Galović Primary School we have been using games in chemistry classes for years as an integral part of teaching. By using directed activities in games, students learn in a fun and creative way the concepts and processes that were previously distant, imaginary, and uninteresting to them, e.g.: the structure of the atom, electron configuration of atoms of various elements, formation of ions, conversion between units of measurement, change of the state of matter, etc. Evaluating the project, we have come to the conclusion that we are on the right track. By introducing games in chemistry classes, we have achieved much better results than in the previous years, all in a fun and motivating way.

REFERENCES

1. V. Velički, T. Topolovčan, *Neuroznanost, nastava, učenje i razvoj govora, Nastava i škola za net-generacije*, Matijević, Milan (ur.), Zagreb: Učiteljski fakultet Sveučilišta u Zagrebu, 2017.
2. E. Jensen, *Poučavanje s mozgom na umu*, Zagreb: Educa, 2005.

FROM OLD TO NEW TOWARDS SUSTAINABLE DEVELOPMENT

Mirta Malčić,^a Suzana Lovrić^b

^a OŠ “Đuro Pilar”, Vinogorska 1, Slavonski Brod, Croatia

^b OŠ “Blaž Tadijanović”, Podvinjska ulica 25, Slavonski Brod, Croatia

mirta.malcic@skole.hr, suzana.lovric1@skole.hr

Recycling is a procedure important for the economy development of every country, for the preservation of the living environment and the sustainability of energy sources. That is why it is crucial to start educating children from a young age, so that as they grow, their awareness of ecology and the importance of recycling will grow as well. Children easily learn and acquire new and useful knowledge on how to live healthier and in harmony with nature. Through the education of the youngest, we achieve a common goal, which is to make environmental protection a priority in the lives of all members of the community.

Waste oils are one of the biggest polluters of our nature, especially water. Most of the waste oil ends up in nature, which poses a danger to the environment: it can make the soil unsuitable for plant growth, and it can have a fatal effect on the animal life in the waters.

The goal of the workshop is to recycle waste oil from fryers by making scented candles. The candles are colored with natural colors (cocoa powder and turmeric) and orange essential oil is added to them. Students prepared for practical work using the teaching method of the flipped classroom.^{1,2}

Flipped classroom is a form of combined learning in which students learn course content online, watching video content at home. After that, they check the adoption of the content with some of the digital tools (Forms)³ and send the output to the teacher who, based on the received answers, plans the lesson at school. At school they discuss and ask the teacher questions about what they watched at home. The teacher intervenes when the students cannot come up with a solution on their own, explaining the problem to them. The goal of the flipped classroom is for students to learn through practical work and by asking questions.

After the finished practical work, the scent of candles enriched with orange essential oil spreads through the classroom.

The adoption of the content and the realization of the outcome are checked with the digital tools *Plickers*⁴ and *Socrative*.⁵

REFERENCES

1. Teachthought, “The definition of the flipped classroom”, [Internet]
2. L. Nielsen, (August 2011) “Five Reasons I’m Not Flipping over the Flipped Classroom. Technology and Learning”, [Internet]
3. <https://e-laboratorij.carnet.hr/google-forms/>
4. <https://e-laboratorij.carnet.hr/plickers/>
5. <https://e-laboratorij.carnet.hr/socrative-kviz-na-drugaciji-nacin/>

CARBONATED WATER: ACID, BUFFER OR BOTH?

Nikola Bedeković

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia

nbedekovic@chem.pmf.hr

When presenting teaching lessons related to buffer solutions, teachers mostly resort to explaining standard examples of acetate or ammonia buffers.^{1,2} Of course, such an approach is not necessarily wrong or bad, but certainly an additional link of the teaching content with everyday substances or phenomena would help the students to learn the material better.

By dissolving pressurized carbon dioxide in water, a chemical reaction occurs between the dissolved gas and water resulting in carbonic acid, commonly known as "carbonated water". Such a solution is acidic with an average pH value of 3.3-4, and could be used as a refreshing drink, or as a base for the production of carbonated juices. It is also frequently used in combination with certain types of wine in noble drink called *gemišt*. In this lecture, the composition of mineral water will be explained on a microscopic level, and its potential as a buffer solution will be determined experimentally.

REFERENCES

1. J. S. Schmuckler, *J. Chem. Educ.* 1982, 59, 305.
2. A. A. Gordus, *J. Chem. Educ.* 1991, 68, 656.

CHALLENGES IN CHEMISTRY CLASSES USING ICT TECHNOLOGY

Monika Pavić, Maja Jurić-Babaja

OŠ Vladimir Nazor, Čepin, Kalnička 17, Croatia

* monika.pavic@skole.hr, maja.juric-babaja@skole.hr

Technology is present in the lives of all of us, including our students. Survey of the population of 7th and 8th grade students shows that students spend four hours online on average. The idea of the activity was to bring chemistry closer to them in an interesting way and so enter their lives.

By correlating chemistry and informatics, students learned to create a website, upload chemistry content to it and organize materials (videos, recordings of experiments, etc.). In this way, children learn and acquire new and useful knowledge more easily, use the Internet, apply learned material and in the end information spreads quickly and is available to everyone at the same time. Students are encouraged to work together, collaborate and work on joint documents for repetition. The goal of this work is to prepare students for life and in this way show them how chemistry and technology go together and complement each other.

We organized the work through joint preparation and cooperation of chemistry and informatics teachers, and through work in chemistry classes (dealing with introductory topics of the 7th grade of elementary school¹). The topics that we put on the website are as follows: Chemistry is a natural science, Trial or experiment, Chemical utensils and accessories, Precautions and protection when performing experiments.²

In the computer science class, students created a website using the Webnode³ online tool. The students were divided into pairs and each pair was given their own section of the website to edit according to the content they received from chemistry. All students worked on one website and were added as page editors. The computer science teacher was the administrator of the website. In the informatics class, the students were introduced to the used tool and its possibilities.

The joint presentation of the work was made through the evaluation of the work according to the given evaluation criteria. We communicated with students via Yammer and checked the adoption of the content and the achievement of the results through Socrativa⁴ i Testmoza.⁵

REFERENCES

1. S. Lukić, I. Marić Zerdum, N. Terenčevska, M. Varga, S. Rupčić Petelinc, Kemija 7, udžbenik kemije u 7. razredu, Školska knjiga
2. <https://edutorij.e-skole.hr/share/proxy/alfresco-noauth/edutorij/api/proxy-guest/985e1c2e-9586-4330-bf1e-c1dc967fdc74/index.html>
3. <https://www.webnode.com/>
4. <https://e-laboratorij.carnet.hr/socrative-kviz-na-drugaciji-nacin/>
5. <https://e-laboratorij.carnet.hr/testmoz-brzi-jednostavni-kvizovi-dostupni-svima/>

HOW TO FACILITATE THE EVALUATION OF STUDENT ACHIEVEMENTS BY USING WRITTEN PAPERS IN CHEMISTRY?

Marijana Bastić

Primary School Rudeš, Zagreb, Croatia

* marijana.bastic@gmail.com

Monitoring and evaluating student achievements is an important part of the curriculum teaching cycle that needs to be carefully planned. In student-oriented teaching for the purpose of adopting educational outcomes, it is possible to achieve it by implementing individual and collaborative project tasks.¹ At the same time, it is necessary to highlight what is expected of the students (goals), how they will work and why (which competencies they will acquire). These competencies are, for example, learning how to learn, learning to search and select information, acquiring communication skills, acquiring skills in the use of ICT and/or laboratory equipment, modeling, creating in some material, etc.

The teacher's autonomy in the development of educational outcomes and evaluation planning enables to achieve certain outcomes in the framework of different thematic units during repetition and practice of the teaching content, some parallel and interdisciplinary, and some through vertical connection.²

Circumstances caused by the Covid-19 pandemic resulted in special conditions for teaching, learning and evaluation, which became predominantly individual in elementary school. Teachers needed to show flexibility both in teaching methods and methods of evaluation. For this purpose, the students' written papers in the form of project tasks that had a given structure proved to be a good choice. The students were introduced in advance to the rubric for evaluating the work, which consisted of the written part and the creation of a model. In addition to the analytical rubric for evaluation, the students were also given a table for self-assessment of the performance of the project task with a rating scale from 2 to 5. At Rudeš Elementary School, seventh-grade students created a brochure about the selected non-metal chemical element at the end of learning about the structure of matter and the periodic table of elements, and eighth graders about the DNA molecule. In addition to their written work, they created a corresponding atom/molecule model. Analysis of the student marks showed that they differ very little from the self-assessment of the quality of the student's work. This method of evaluation contributes to the satisfaction of the students³ through the process of checking knowledge and the final evaluation, and makes it easier for the teacher to assess the quality of the students' knowledge.

REFERENCES

1. M. Matijević, D. Radovanović, *Nastava usmjerena na učenika*, Zagreb: Školske novine, **2011**.
2. https://narodne-novine.nn.hr/clanci/sluzbeni/2019_01_10_208.html (Odluka o donošenju kurikuluma za nastavni predmet Kemije za osnovne škole i gimnazije u Republici Hrvatskoj), 7.10.2022.
3. E. Jensen, *Super-nastava*, Zagreb, Educa, **2005**.

FOSTERING TRANSDISCIPLINARY LEARNING IN SCIENCE AND TECHNOLOGY

Jurica Bauer

MERLN Institute for Technology-Inspired Regeneration, Faculty of Health, Medicine and Life Sciences,
Maastricht University, Universiteitssingel 40, 6229 ER Maastricht, The Netherlands

j.bauer@maastrichtuniversity.nl

The presented work will highlight two education initiatives from Dutch universities at the undergraduate level. These examples reflect and illustrate the recent trends in science and technology, as well as in the corresponding education programs and processes, to work and study not only at the interface of various disciplines but also across disciplines in order to achieve creative solutions to ongoing societal challenges.

The first initiative concerns an innovative approach to teaching nanotechnology as part of the undergraduate Chemistry curriculum at the Inholland University of Applied Sciences in Amsterdam¹. Students work in pairs to review a relevant and contemporary multidisciplinary topic in nanotechnology and to propose and develop an original research idea. This is presented and defended in the form of an oral presentation before an audience of peers and is written using an article template. The aim of the described approach is to not only help students learn about nanotechnology in a fun and active manner but also to stimulate critical scientific thinking in students as well as to foster the development of their academic, professional, and soft skills. In line with this, the assessment of students relies on the assessment of the oral presentation, engagement in discussion, and the written research proposal.

The second initiative comes from the Maastricht University where a multidisciplinary Bachelor program with the focus on Regenerative Medicine & Technology is being set up. Regenerative Medicine is a relatively new field found at the intersection of science, engineering and medicine. Applying the principles of problem- and research-based learning the new program intends to offer a solid science and engineering foundation strongly integrated with the relevant aspects of medicine and entrepreneurship. As such, the program aims to educate a new generation of researchers that will be able to swiftly adapt to any area of regenerative medicine, and contribute to the design and development of medical therapies, products and devices for research and clinical use. The desired competencies and final qualifications have been defined and translated into a concept curriculum. The education is now being developed in further detail keeping the principles of constructive alignment and the CCCS (Constructive, Contextual, Collaborative, Self-Directed) learning principles in mind. The program is undergoing the initial accreditation process at the moment and is intended to start in September 2023 pending a positive decision by the accreditation committee.

ACKNOWLEDGEMENTS The Department of Life Sciences and Chemistry at the Inholland University of Applied Sciences, including their Chemistry student population, is acknowledged for the support and freedom to design the Nanotechnology course in its described form. The Faculty of Health, Medicine and Life Sciences at the Maastricht University and especially the MERLN Technology-Inspired Institute for Regenerative Medicine are acknowledged for the opportunity to lead the design and development of the new Bachelor program Regenerative Medicine & Technology as well as all the support during this project.

REFERENCES

1. J. Bauer, *J. Chem. Educ.* **2021**, *98*, 2347

THE CURRENT CHALLENGES OF IMPROVING CHEMISTRY LEARNING

Olgica Martinis

Education and Teacher Training Agency, Donje Svetice 38, Zagreb, Croatia

olgica.martinis@azoo.hr

We face numerous and different challenges in day-to-day life. New challenges in teaching the chemistry have caused the changes in the existing way of teaching it.¹ In order to discuss successfully possible solutions for improving the teaching of chemistry, it is necessary to understand the current encounters. One of the current challenges of teaching chemistry are changes in teaching strategies in certain periods. Participants will learn about the dominant teaching strategies during chemistry classes at the beginning of the 20th century. Therefore, we will talk about the behaviorist theory of teaching and the application of the cognitivist theory that applied in most of Europe, as well as the increasingly widespread use of the constructivist theory.² By applying this theory, the student is placed in the center of the teaching process, and thus the teaching performance is different. Student's activities are planned and based on their own experience during chemistry classes, especially when guided teaching is applied. Furthermore, the strategy of independent teaching can be applied to the activities for gifted students. The need for an appropriate choice and elaboration of a teaching strategy depending on the predictions and needs of the students and on the teaching situation is evident.³ Since teaching strategies are part of the curriculum circle, the participants will be able to conclude about the need to change each component of the curriculum circle, if there is a change in only one of them. Application of experiential teaching is desirable and possible in the conditions of face-to-face and online teaching. It is essential to plan the activities of students who will have their own experience, regardless of the way of teaching (face-to-face or online) and the form of work (individual, in pairs or groups).⁴ One of the competencies of a successful teacher refers to a teacher who knows how to recognize successfully conducted lessons and what and at what point has to be changed in order to make chemistry lessons more successful.⁵

Successful chemistry teaching is the result of the synergy of the work of students and teachers, who understand the professional content, apply a pedagogical approach and regularly evaluate their work, regardless of current challenges regarding the application of teaching strategies and other factors that affect the quality of chemistry teaching.

REFERENCES

1. J. D. Herron, S. C. Nurrenbern, *J. Chem. Edu.* **1999**, *76*, 1353.
2. M. Matijević, D. Radovanović, *Nastava usmjerena na učenika, Školske novine*, **2011**, Zagreb.
3. B. Bognar, *Croat. J. Educ.* **2016**, *18*, 225.
4. E. Jensen, I. Marušić, N. M. Šoljan, *Super-nastava: nastavne strategije za kvalitetnu školu i uspješno učenje*, **2003**, Educa.
5. V. Makarevics, *J. Teach. Educ. Sustain.* **2008**, *10*, 68.

POSTER PRESENTATIONS

SELECTIVE GUEST BINDING AND RELEASE: TUNING THE PROPERTIES OF OCTAMERIC NANOCAPSULES


Wiktorija Adamska,^{a,*} Grzegorz Markiewicz,^a Artur R. Stefankiewicz^{a,b}

^a Faculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland, ^b Center for Advanced Technology, Adam Mickiewicz University, Uniwersytetu Poznańskiego 10, 61-614 Poznań, Poland

* wikada2@amu.edu.pl

Supramolecular chemistry is built on non-covalent interactions including hydrogen bonds, π - π interactions, and van der Waals forces. By controlling molecular self-assembly and recognition processes through the application of these weak intermolecular forces, it is feasible to produce complex supra-structures from synthetically simple molecules that have adjustable functions and features. Many other self-assembled supramolecular structures, such as capsules, have been discovered throughout the years.¹ These structures are of great interest to many scientists because of their singular capacity to attach the guest molecules inside clearly defined internal spaces.

Our team presented the first hydrogen-bonded octameric capsule based on the benzene-1,3,5-tricarboxylic acid (BTA) core a few years ago.² The capsule is held together by 48 cooperative hydrogen bonds. Thanks to a large internal cavity of 1719 Å³, this assembly is an excellent receptor for C70 and C60 fullerenes, with encapsulation of C70 being highly favored. In our recent research, we outline a synthetic approach to finely tune and target the solubility of the octameric capsules for preparative guest separation in addition to the encapsulation of the catalytically-active species.



ACKNOWLEDGEMENTS This work was supported by the National Science Centre (Grant: SONATA BIS 2018/30/E/ST5/00032)

REFERENCES

1. L. Adriaenssens, P. Ballester, *Chem. Soc. Rev.* 2013, 42, 3261.
2. G. Markiewicz, A. Jenczak, M. Kołodziejcki, J. J. Holstein, J. K. M. Sanders, A. R. Stefankiewicz, *Nat. Commun.* 2017, 8, 15109.

STRUCTURAL AND THERMODYNAMIC FEATURES OF DIAMONDROID AMMONIUM SALT INCLUSION COMPLEXES WITH CYCLODEXTRINS

Marija Alešković,^{a,*} Amalija Ernečić,^a Sunčica Roca,^b Viktoriia Lazarenko,^c
Nikola Bregović,^c Marina Šekutor^a

^a Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička c. 54, Zagreb, Croatia, ^b NMR Center, Ruđer Bošković Institute, Bijenička c. 54, Zagreb, Croatia,


^c Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* marija.aleskovic@irb.hr

Functionalized diamondoid ammonium salts are an intriguing class of compounds capable of forming ultra-stable complexes with cucurbituril hosts (CB[n]) in aqueous environment with binding constants up to $10^{15} \text{ mol}^{-1} \text{ dm}^3$.¹ Inspired by these properties, we decided to broaden the research scope of inclusion complexes stabilized by hydrophobic interactions between various cage ammonium salt guests and cyclodextrin (CD) as a different macrocycle host molecule.²

Herein we present the synthesis of a continued series of diamondoid ammonium salts and a detailed thermodynamic study of their complexation with β - and γ -CDs. The complex stability constants were determined by ITC and ^1H NMR titrations in aqueous solutions. Additional NMR spectroscopic techniques (^1H - ^1H NOESY, ^1H DOSY) along with computational methods were employed to provide a deeper insight into the thermodynamics of hydrophobically driven complexations and structural features of the formed supramolecules.

The binding strength was assessed as a function of diamondoid salt size, functional group position and cyclodextrin cavity size and by varying these parameters, a structure-selectivity relationship can be determined in order to facilitate the design of next generation of suitable guests.



ACKNOWLEDGEMENTS This work has been supported by the Croatian Academy of Science and Arts and the Croatian Science Foundation (UIP-2017-05-9653 (DiamMat), IP-2019-04-9560 (MacroSol)).

REFERENCES

1. D. Sigwalt, M. Šekutor, L. Cao, P. Y. Zavalij, J. Hostaš, H. Ajani, P. Hobza, K. Mlinarić-Majerski, R. Glaser, L. Isaacs, *J. Am. Chem. Soc.* 2017, *139*, 3249.
2. M. Alešković, S. Roca, R. Jozepović, N. Bregović, M. Šekutor, *New J. Chem.* 2022, *46*, 13406.


SUSTAINABLE SOLUTION FOR SYNTHESIS OF DIAMONDROID ETHERS

Jasna Alić Stolar,^{a,*} Tomislav Stolar,^b Zoran Štefanić,^b Krunoslav Užarević,^b Marina Šekutor^a

Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia, ^b Division of Physical Chemistry, Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia

* jasna.alic@irb.hr

Diamondoids are diamond-like hydrocarbon cage molecules that have emerged as promising candidates in a wide range of applications.^{1, 2} Their inherent properties and availability for selective functionalization allows for a development of variously applicable scaffolds. However, the tendency of smaller diamondoid derivatives to sublime at even slightly elevated temperatures coupled with solubility issues of larger derivatives offers a challenge for their preparation. Here, we present the first mechanochemical synthesis of several diamondoid ethers differing in the size and number of their cage subunits. Efficient mechanochemical preparation of such ethers is enabled solely by high-temperature ball milling conditions³ and does not proceed under ambient conditions. When compared to the conventional synthesis⁴ of the same ether derivatives, the calculated green chemistry metrics showed significant sustainability benefits of the mechanochemical approach. Thus, mechanochemical procedures are faster, allow for the use of an inorganic green base, result in comparable or superior reaction yields, and are overall more sustainable and eco-friendlier. Furthermore, application of mechanochemistry in the preparation of large diamondoid composites opens up a sustainable approach to the synthesis of next-generation materials consisting of diamondoid subunits, unhampered with solubility capabilities of the reactants.



ACKNOWLEDGEMENTS We acknowledge financial support from the Croatian Science Foundation (UIP-2017-05-9653 and IP-2020-02-4702).

REFERENCES

- H. Schwertfeger, A. A. Fokin, P. R. Schreiner, *Angew. Chem. Int. Ed.* 2008, 47, 1022; M. A. Gunawan, J.-C. Hierso, D. Poinso, A. A. Fokin, N. A. Fokina, B. A. Tkachenko, P. R. Schreiner, *New J. Chem.* 2014, 38, 28.
- G. C. Mcintosh, M. Yoon, S. Berber, D. Tomanek, *Phys. Rev. B.* 2004, 70, 045401.
- N. Cindro, M. Tireli, B. Karadeniz, T. Mrla, K. Užarević, *ACS Sustainable Chem. Eng.* 2019, 7, 16301.
- J. Alić, I. Biljan, Z. Štefanić, M. Šekutor, *Nanotechnology* 2022, 33, 355603.

COMPLEXATION THERMODYNAMICS OF BENZENE DERIVATIVES WITH MACROCYCLIC RECEPTORS

Tea Babić,* Andrea Usenik, Katarina Pičuljan, Nikola Bedeković, Josip Požar

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia

* tea.babic@chem.pmf.hr

Cucurbit[7]uril (**CB7**) and β -cyclodextrin (**β -CD**) are well-known macrocyclic receptors for hydrophobic compounds in aqueous solutions.^{1,2} Since the host and guest engage in relatively weak van der Waals interactions, the stability of formed complexes is predominantly ascribed to the thermodynamically favourable dehydration of the host cavity³ and the guest.⁴ In contrast with **β -CD**, the inclusion of simple benzene derivatives within **CB7** was not systematically explored. A variety of mono- and 1,4-disubstituted benzene derivatives with different electron-donor (-acceptor) properties were hence chosen as potential guests for heptameric glycouril receptor. The corresponding complexation reactions were explored by means of isothermal titration microcalorimetry and ROESY NMR spectroscopy in a wide temperature range. Guests functionalized with electron-donor groups formed more stable complexes with **CB7** due to more favourable complexation energetics. The stability of **β -CD** complexes was always lower than that of the corresponding **CB7** complexes because of the less favourable enthalpic contribution to $\Delta_r G^\circ$. Significant temperature dependence of $\Delta_r H^\circ$ and $\Delta_r S^\circ$ resulted in an almost complete enthalpy-entropy compensation. These results suggest that the guests behave as structure makers at lower temperatures (classic iceberg hydration model) and as structure breakers at higher explored temperatures.⁴




Figure 1. Schematic representation of the complexation process.

ACKNOWLEDGEMENTS This research was fully supported by the Croatian Science Foundation (project MacroSol, IP-2019-04-9560) and European Regional Development Fund (project CluK, KK.01.1.1.02.0016).

REFERENCES

1. S. J. Barrow, S. Kasera, M. J. Rowland, J. del Barrio, O. A. Scherman, *Chem. Rev.* **2015**, *115*, 12320.
2. M. V. Rekharsky, Y. Inoue, *Chem Rev.* **1998**, *98*, 1875.
3. F. Biedermann, W. M. Nau, H.-J. Schneider, *Angew. Chem. Int. Ed.* **2014**, *53*, 2.
4. K. Leko, M. Hanževački, Z. Brkljača, K. Pičuljan, R. Ribić, J. Požar, *Chem. Eur. J.* **2020**, *26*, 5208.


STOICHIOMETRIC DIVERSITY OF HALOGEN BONDED COCRYSTALS DERIVED FROM *o*-DIIODOTETRAFLUOROBENZENE AND MONOTOPIC ACCEPTORS

Nikola Bedeković, Vladimir Stilinović, Dominik Cinčić

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia

* nbedekovic@chem.pmf.hr

Perfluorinated halogenobenzenes are a group of halogen bond donors that include a wide range of compounds with different number of donor atoms (i.e. different topicities), what ultimately allows to synthesize halogen bonded cocrystals with various stoichiometries and crystal packing motifs.^{1,2} The structural and the statistical study of the compounds including 1,3- and 1,4-diiidotetrafluorobenzene and simple monotopic acceptors have pointed out significant differences in their topicities in corresponding crystal structures: **14ditfb** has been found to be mostly ditopic, while **13ditfb** is generally monotopic halogen bond donor.³ The third isomer – **12tfib** – consists of two donor atoms located at an angle of 60° one to another, which can be a limiting factor for the formation of two halogen bonds simultaneously, due to unfavourable steric influences between the acceptor molecules. To test this hypothesis, we have prepared and structurally characterized six binary cocrystals of **12tfib** and six simple monotopic pyridines (3-picoline, 2,6-lutidine, 3,4-lutidine, 3,5-lutidine, 2,4,6-collidine and isoquinoline). Four obtained cocrystals are of 1:1, and two are of 1:2 stoichiometric ratio, while **12tfib** has served as a ditopic donor in total four structures (two with 1:1 and two with 1:2 stoichiometry). In 1:1 cocrystals with 3-picoline and isoquinoline acceptor molecules participate in bifurcated halogen bonds, while enabling ditopicity of the **12tfib** molecule in those compounds. Halogen bonded complexes in 1:2 cocrystals are not planar, but were found in bent conformations due to the unfavourable steric repulsions of two halogen-bonded acceptor molecules.



ACKNOWLEDGEMENTS This research was supported by the Croatian Science Foundation under the project IP-2019-04-1868.

REFERENCES

1. N. Bedeković, V. Stilinović, T. Friščić, D. Cinčić *New J. Chem.* 2018, 42, 10584.
2. P. Metrangolo, G. Resnati, *Angew. Chem. Int. Ed.* 2000, 39, 1782.
3. N. Bedeković, T. Piteša, M. Eraković, V. Stilinović, D. Cinčić. *Cryst. Growth Des.* 2022, 22, 2644.

MOLECULAR MODELLING OF MntR TRANSCRIPTION FACTOR FROM *BACILLUS SUBTILIS*


Branimir Bertoša,* Zoe Jelić Matošević

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* branimir.bertosa@chem.pmf.hr

Manganese homeostasis is a highly sensitive process in bacteria. On the one hand, manganese ions are essential for the survival of a bacterium since they are necessary for basic cellular processes, such as DNA replication and resistance to oxidative stress. At the same time, manganese ions are toxic for the cell if present in high concentrations. Therefore, a bacterium must have a sensitive cellular mechanisms for regulation of manganese homeostasis.¹ In case of the bacterium *Bacillus subtilis* this mechanism relies on the MntR protein which is a transcriptional repressor that is activated for DNA binding by manganese ions (Mn^{2+}).¹ Although extensive experimental data on *B. subtilis* MntR is available, the molecular mechanism through which *B. subtilis* MntR is activated for interaction with DNA upon Mn^{2+} binding is still a puzzle.

Starting from available crystal structures of different forms of this protein, systems with and without Mn^{2+} bound in the binding sites were prepared for molecular dynamics (MD) simulations. Special attention was put on protein protonation and different approaches for adding polar hydrogen atoms were applied since hydrogen bond networks might have an important role in the protein's molecular mechanisms. The results of analyses of all-atom MD simulations point to the changes in the protein's dynamics as the most pronounced change due to Mn^{2+} binding. It seems that Mn^{2+} binding reduces the conformational space of the protein and "locks" the DNA binding helices in the conformation appropriate for DNA binding. In order to validate this hypothesis, mutations that should induce activity of *B. subtilis* MntR even in the absence of Mn^{2+} ions were constructed *in silico* and computationally studied.



ACKNOWLEDGEMENTS This work was supported by the Croatian Science Foundation project "Manganese metallosensors" IP-2020-02-3446.

REFERENCES

1. P. Chandrangsu, C. Rensing, J. D. Helmann, *Nat. Rev. Microbiol.* **2017**, *15*, 338.

BIOLOGICAL ACTIVITY OF 3-AMINOQUINUCLIDINE QUARTERNARY SALT

Katijana Bilandžić,* Renata Odžak, Matilda Šprung

Department of Chemistry, Faculty of Science, R. Bošković 33, 21000 Split, Croatia

* kbilandzi@pmfst.hr

Quaternary ammonium compounds (QACs) are amphiphilic compounds with exceptional properties and a wide range of applications. They display very good biological activities. Due to the development of bacterial resistance to QACs, the current aim is to synthesize new QACs which are biodegradable, but still efficient.¹ Quaternization of heterocyclic quinuclidine and its derivatives has proven to be a good solution since its quaternary salts have great antibacterial and antifungal properties. For this research, the antibacterial activity of synthesized 3-aminoquinuclidine quaternary salt with an alkyl chain of 16 carbon atoms has been investigated by microdilution method on seven bacterial strains: *Staphylococcus aureus*, *Escherichia coli*, *Pseudomonas aeruginosa*, *Enterococcus faecalis*, *Salmonella enteritidis*, *Listeria monocytogenes* and *Bacillus cereus*. The best MIC value obtained was 9 nM against two Gram-positive bacteria: *L. monocytogenes* ATCC 7644 and *S. aureus* ATCC 25923. The second-best MIC was against Gram-positive *E. faecalis* ATCC 29212 at 36 nM. Obtained values are considered the potential for future investigations of that quaternary ammonium salt.

REFERENCES

1. M. C. Jennings, K. P. C. Minbiole, W. M. Wuest, *ACS Infect. Dis.* **2016**, *1*, 288.


PREPARATION AND MASS SPECTRAL CHARACTERIZATION OF DIAMONDROID DERIVATIVES

Nataša Burić,* Marina Šekutor

Department of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia

* nburic@irb.hr

Diamondoids are hydrocarbons with characteristic cage structures¹ and in nature these molecules are predominately found in mature oils as final products of fossil decomposition, indicating their high thermodynamical stability.² They were named after diamond, but unlike diamond diamondoids can be selectively functionalized since they possess terminal hydrogen atoms with different reactivity. Diamondoid derivatives have already found application in many fields,² with preparation of new molecules constantly in progress. However, derivatization of diamondoid cages can often lead to formation of several products and many of them are constitutional isomers. Such isomers can show different fragmentation pathways in mass spectrometry, making gas chromatography with mass spectrometric detection (GC-MS) an ideal tool for monitoring the reactions, analyzing the reaction mixtures and characterizing the isolated products.³ For example, a common route in diamantane cage functionalization is its direct



oxidation,⁴ a reaction that benefits from usage of the GC-MS analytics since slight variations in the reaction conditions result in the formation of different products in different ratios (Figure 1). Here we will showcase the application of combined chromatographic and mass spectral characterization for fast reaction screening.

Figure 1. Oxidation of diamantane leads to different products.

ACKNOWLEDGEMENTS

This work was supported by the Croatian Science Foundation (UIP-2017-05-9653).

REFERENCES

1. H. Schwertfeger, A. A. Fokin, P. R. Schreiner, *Angew. Chem. Int. Ed.* **2008**, *47*, 1022; M. A. Gunawan, J.-C. Hierso, D. Poinso, A. A. Fokin, N. A. Fokina, B. A. Tkachenko, P. R. Schreiner, *New J. Chem.* **2014**, *38*, 28.
2. J. E. Dahl, S. G. Liu, R. M. K. Carlson, *Science* **2003**, *299*, 96.
3. L. Vodička, J. Burkhard, J. Janků, *Collect. Czech. Chem. Commun.* **1986**, *51*, 1086.
4. A. Fokina, Boryslav A. Tkachenko, A. Merz, M. Serafin, J. E. P. Dahl, R. M. K. Carlson, A. A. Fokin, P. R. Schreiner, *Eur. J. Org. Chem.* **2007**, *28*, 4738.

THERMODYNAMICS OF SEVERAL ION-PAIRINGS IN ACETONITRILE

Marija Cvetnić,* Davor Mendeš, Nikola Bregović, Vladislav Tomišić

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia

* marija.cvetnic@chem.pmf.hr

In the field of supramolecular chemistry, ditopic receptors capable of binding ion-pairs play a significant role.¹ The binding of cation on a ditopic receptor often allosterically affects the binding of an anion (or *vice versa*) enhancing the stability of the resulting complexes. In order to fully understand and exploit such cooperativity it is necessary to characterize the underlying structural and thermodynamic aspects.²

Recently we synthesized amide-ureido-calix[4]arene derivative and investigated its ability as ion-pair receptor in acetonitrile. During this investigation a need for the values of equilibrium constants for ion-pairing in this solvent emerged. Namely, organic solvents are generally better media for the complexation of ions with organic receptors because ions are poorly solvated in them in comparison to water. However, the latter fact also results with the increase of significance of ion-pairing in organic solvents. As the literature lacks the constants of ion-pairings in acetonitrile, we investigated the thermodynamics of ion-pairing for several lithium and sodium salts using conductometric and microcalorimetric titrations. The obtained results were correlated with the basicities of the examined anions in acetonitrile,^{3,4} and compared with standard Gibbs energies of transfer of the appropriate ions from water to acetonitrile.⁵

ACKNOWLEDGEMENTS This work has been fully supported by the Croatian Science Foundation under the project IP-2019-04-9560 (MacroSol).

REFERENCES

1. Q. He, G. I. Vargas-Zúñiga, S. H. Kim, S. K. Kim, J. L. Sessler, *Chem. Rev.* **2019**, *119*, 9753.
2. B. Qiao, A. Sengupta, Y. Liu, K. P. McDonald, M. Pink, J. R. Anderson, K. Raghavachari, A. H. Flood, *J. Am. Chem. Soc.* **2015**, *137*, 9746.
3. A. Kütt, S. Tshepelevitsh, J. Saame, M. Lõkov, I. Kaljurand, S. Selberg, I. Leito, *Eur. J. Org. Chem.* **2021**, 1407.
4. D. Barišić, V. Tomišić, N. Bregović, *Anal. Chim. Acta* **2019**, *1046*, 77.
5. Y. Marcus, *Ion Properties*, Taylor & Francis Inc, Boca Rosa (USA), **1997**.

RHEOLOGICAL CHARACTERIZATION – IMPORTANT TOOL IN REVERSE ENGINEERING OF OPHTHALMIC SOLUTIONS

Karla Dubaić,* Leo Štefan, Maša Safundžić Kučuk

Jadran galenski laboratorij d.d. (JGL d.d) Svilno 20, 51000, Rijeka, Hrvatska

* karla.dubaic@jgl.hr

Eye drops are the prevalent ophthalmic drug delivery system with certain limitations in terms of delivery to ocular tissues. To improve delivery, polymers are often added in formulations due to their unique properties such as prolonged release and improved drug solubility.¹ Development of the topical generic ophthalmic drug products revolves around demonstration of the equivalence to the reference medicinal product (RMP). If RMPs contain polymers, reverse engineering studies are conducted to determine grade and concentration of added polymer, to ensure required similarity. Due to complexity of the polymers, and their non-Newtonian behavior, several analytical techniques are applied during this process, however rheological characterization is essential for the final confirmation of equivalence with RMP² because it can be used to compare the behavior of the products on the eye. Tests that are required are: shear rate sweep test (viscosity curve), amplitude sweep and frequency sweep. These tests gather information about visco-elastic properties of added polymer in formulation. For more complex ophthalmic topical systems (e.g., in situ gels), gelation time of polymer with tear fluid provides information about gel characteristics of the final product, and its behavior during application. Presented work provides overview of different rheological tests and points out their importance during development of ophthalmic solutions.




Figure 1. Viscosity curves obtained for JGL formulation (blue) and RMP (green) containing the same type of thickener, but different grade based on molecular weight (MW). Although the same at one point (arrow), during applied shear rate range, viscosity profile is different.

REFERENCES

1. M. M. Allyn, R. H. Luo, *Front. Med.* **2022**, *8*, 787644.
2. A. Simoes, M. Miranda, *Pharmaceutics* **2020**, *12*, 820.

CALCIUM PHOSPHATE/TITANATE NANOMATERIAL COMPOSITES FOR BIOMEDICAL APPLICATIONS

Ina Erceg,^{a,*} Marija Milić,^b Krunoslav Ilić,^c Darija Domazet Jurašin,^a Nadica Maltar Strmečki,^d
Ivana Vinković Vrček,^c Maja Dutour Sikirić^a

^a Laboratory for Biocolloids and Surface Chemistry, Division of Physical Chemistry,
Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia, ^b Department of Clinical Laboratory
Diagnostics, University Hospital Centre Osijek, Huttlerova 4, Osijek, Croatia,

^c Institute for Medical Research and Occupational Health, Ksaverska cesta 2, Zagreb, 10000, Croatia, ^d
Laboratory for Electron Spin Spectroscopy, Division of Physical Chemistry, Ruđer Bošković Institute,
Bijenička c. 54, 10000 Zagreb, Croatia

* ierceg@irb.hr

In recent years, calcium phosphates (CaPs) composites with TiO₂ nanomaterials (TiNMs) have emerged as a promising solution to the poor mechanical properties of CaPs based bone regeneration materials.¹ Using TiNMs of different morphologies opens up an opportunity to fine-tune the properties of composites. Among different TiNMs, the formation of CaPs has been investigated most frequently on nanotube arrays, while the studies on other TiNMs are scarce. To fill this gap, in this study CaPs composites with TiNMs of different structures, compositions, and morphologies were prepared by precipitation at room temperature.^{2,3} Four different TiNMs were used: anatase nanoparticles and nanoplates, titanate nanotubes and TiO₂ (B) / hydrogen titanate nanowires. In all cases, calcium-deficient apatite (CaDHA) was formed after one hour of aging. Although the morphology and size of the crystalline domains of CaDHA obtained on the TiNMs were different, no significant difference in their local structure was observed. To test the potential of the composites for biomedical applications, their effect on hemostasis, cell viability of Jurkat T-cells and cytokine expression in Jurkat T-cells were investigated. The largest effect on hemostasis and cytokine expression, as well as a significant reduction of Jurkat cell viability after 24 hours was observed for composites with nanotubes and nanowires.

The obtained results indicate a rapid and versatile method for the preparation of CaP/TiNMs composites and confirm the potential of such composites for biomedical applications.

ACKNOWLEDGEMENTS Financial support from the Croatian Science Foundation, Grant HRZZ-IP-2018-01-1493 is greatly acknowledged.

REFERENCES

1. A. K. Vemulapalli, R. M. R. Penmetsa, R. Nallu, R. Sireyala, *J. Compos. Mater.* **2020**, *54*, 765.
2. I. Erceg, A. Selmani, A. Gajović, I. Panžić, D. Iveković, F. Faraguna, S. Šegota, M. Čurlin, V. Strasser, J. Kontrec, D. Kralj, N. Maltar Strmečki, M. Dutour Sikirić, *Colloids Surf., A* **2020**, *593*, 124615.
3. I. Erceg, A. Selmani, A. Gajović, B. Radatović, S. Šegota, M. Čurlin, V. Strasser, J. Kontrec, D. Kralj, N. Maltar-Strmečki, R. Barbir, B. Pem, I. Vinković Vrček, M. Dutour Sikirić, *Nanomaterials*, **2021**, *11*, 1523.

CAN HALOGEN BOND PAY THE ENTROPY PENALTY?

Luka Fotović, Nikola Bedeković, Josip Požar, Vladimir Stilinović

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* lfotovic@chem.pmf.hr

Halogen bond is an attractive, strong and directional noncovalent interaction in which a halogen atom acts as a Lewis acid.¹ Most commonly used acceptors of halogen bonds are anions² and neutral molecules containing nitrogen or oxygen atoms³ – i.e., species capable of acting as Lewis bases. Typical halogen bond donors are perfluorinated iodobenzenes,^{4,5} whereas the donor properties of perfluorinated iodoalkanes, such as 1, 4-diiodooctafluorobutane (**ofib**) remained almost completely unexplored.

This type of bonding has attracted considerable attention in the solid state chemistry, where it is widely used for the synthesis of halogen bonded cocrystals. According to the most recent definition, cocrystals are solids that are crystalline materials composed of two or more molecules/coformers in the same crystal lattice.⁶ Cocrystals are commonly composed of two conformers, which as separate substances are solids at room temperature. Less commonly one conformer will be a liquid (unless it was also the crystallisation solvent and therefore present in huge excess), while cases where two liquids at room temperature produce a solid cocrystal are extremely rare, as formation of a solid from two liquids is extremely entropically unfavourable.⁷

This study reports the structures of five new cocrystals of the aliphatic ditopic **ofib** and monotopic nitrogen-based aromatic halogen bond acceptors which were obtained by reaction of liquid halogen bond donor (**ofib**) and liquid acceptor in 1:2 molar ratio. Single crystal X-ray diffraction analysis of obtained crystal products revealed that the cocrystals are comprised of discrete C–I⋯N halogen bonded complexes of 1:2 stoichiometry. The $\Delta_r H$ values for the reaction of **ofib** and investigated acceptors were determined calorimetrically at 25 °C. As expected, the cocrystal formation was strongly exothermic in all cases. The particularly favourable halogen bonding realized within the products can thus overcome the entropy penalty arising from association of two liquid reactants into a solid at room temperature.

ACKNOWLEDGEMENTS. This work has been supported by the Croatian Science Foundation under the project IP-2019-04-1868.

REFERENCES

1. P. Metrangolo, G. Resnati, *Chem. Eur. J.* 2021, 7, 2511.
2. L. Fotović, V. Stilinović, *CrystEngComm* 2020, 22, 4039.
3. N. Bedeković, L. Fotović, V. Stilinović, D. Cinčić, *Cryst. Growth Des.* 2022, 2, 987.
4. N. Bedeković, V. Stilinović, T. Friščić, D. Cinčić, *New J. Chem.* 2018, 42, 10584.
5. N. Bedeković, T. Piteša, M. Eraković, V. Stilinović, D. Cinčić, *Cryst. Growth Des.* 2022, 4, 2644.
6. M. J. Zaworotko *et al.*, *Cryst. Growth Des.* 2012, 5, 2147.
7. A. J. Cruz-Cabeza, S. E. Wright, A. Bacchi, *Chem. Commun.* 2020, 56, 5127.

LC-MS/MS AS A SOLUTION IN PRECISE, SELECTIVE, CONFIRMATORY AND COST EFFECTIVE FOOD MULTI-CONTAMINANT ANALYSIS

Josipa Golenja,^{a,*} Michael Sulyok,^b Ivana Dodlek Šarkanj,^a Bojan Šarkanj^a

^a Department of Food Technology, University North, Trg dr. Žarka Dolinara 1, HR-48000 Koprivnica, Croatia, ^b Institute of Bioanalytics and Agro-Metabolomics, Department of Agrobiotechnology (IFA-Tulln), University of Natural Resources and Life Sciences, Vienna, Konrad Lorenzstr. 20, A-3430 Tulln, Austria

* jogolenja@unin.hr

The LC-MS/MS has over the last decade matured into the leading technique for detection of organic contaminants in Food and Feed. The sensitivity, selectivity and robustness of LC-MS/MS instruments has continuously improved, positioning them in legislation as confirmatory instruments. Although initially it was not possible to obtain accurate results down to the ppb range (as required in legislation) without any matrix removal and analyte enrichment, the latest generation of instruments can use dilute & shoot approach. This is one of the pre-requisites for multi-contaminant analysis due to widely different physico-chemical properties of all organic contaminants that are legally regulated.

The scheduled MRM (multiple reaction monitoring) mode has enabled acquisition of large analyte lists, e.g., multimycotoxins, bacterial metabolites, veterinary drug residues, pesticide residues, where methods covering more than 500 compounds have become fairly routine.¹

We have applied a LC-MS/MS multicontaminant analysis method covering more than 500 fungal metabolites and dilute & shoot approach for analysis of multiple legally regulated mycotoxins with indicative levels and unregulated mycotoxins in cereals and feed samples from Croatia.

The results showed that multiple contaminants are predominantly present in analyzed cereals and feed, usually grouping within one fungal genera metabolites. The 100 % co-occurrence of mycotoxins with some bacterial and yeast metabolites was confirmed and the usage of such dilute & shoot with LC-MS/MS multi-contaminant method was justified.

ACKNOWLEDGEMENTS The author would like to thank the VIP Beer project for support. Josipa Golenja would like to thank University North for support, as well as her family and friends.

REFERENCES

1. M. Sulyok, D. Stadler, D. Steiner, R. Krska, *Anal. Bioanal. Chem.* **2020**, *412*, 2607.

STRUCTURE COMPARISON AND STRUCTURAL ALIGNMENTS OF OLIGOMERIC PURINE NUCLEOSIDE PHOSPHORYLASES

Boris Gomaz, Zoran Štefanić

Division of Physical Chemistry, Ruđer Bošković Institution, Bijenička cesta 54, Zagreb, Croatia

* bgomaz@irb.hr

It is known that enzymes can be composed of one or more, same or different, subunits. If some protein is composed of more than one subunit, it is called an oligomeric protein,¹ and one example of such proteins are purine nucleoside phosphorylases (PNPs). There are two main classes of this enzyme: trimeric PNP or “low molecular mass” protein which can mostly be found in eukaryotic organisms, and hexameric PNP or “high molecular mass” protein which can mostly be found in prokaryotic organisms (Figure 1).² These two types of enzymes share only 20 - 30 % sequence identity, but the overall fold of the single monomer is similar, and yet this similar monomeric building block makes a different quaternary structure. A large number of 3D structures of PNPs have been determined so far. We can base our understanding of the mechanism by which monomeric subunits communicate in these enzymes on this structural data, and take into account the oligomeric arrangement of subunits in enzymes from different organisms.

As part of the ALOKOMP project (alokomp.irb.hr) which tries to understand the allosteric communication between these monomeric subunits, the relational database of PNPs is constructed. As a special subset of this database, an all-to-all overlap of these structures is also made, where we can cluster this data as well as compare some of the structures according to their assembly symmetry and their crystallographic symmetry.




Figure 1. Monomeric units from trimeric PNP and hexameric PNP show similar overall fold but form different quaternary structures.

ACKNOWLEDGEMENT This work was funded by the Croatian Science Foundation, Grant number IP-2019-04-6764.

REFERENCES

1. B. Kojić-Prodić, Z. Štefanić, *Symmetry* **2010**, *2*, 884.
2. M. Narczyk, B. Bertoša, L. Papa, V. Vuković, I. Lešćić Ašler, B. Wielgus-Kutrowska, A. Bzowska, M. Luić, Z. Štefanić, *FEBS J.* **2018**, *285*, 1305.

A COMPUTATIONAL STUDY OF CALIXARENE-CYCLOPEPTIDE ANION RECEPTORS

Gordan Horvat,^a Filip Kučas,^a Neven Golenić^b

^a Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia, ^b Department of Physics, Faculty of Science, University of Zagreb, Bijenička c. 32, Zagreb, Croatia

* ghorvat@chem.pmf.hr

Calixarenes are macrocyclic oligomers consisting of phenolic subunits linked by methylene bridges in the *ortho*-position. Unsubstituted calixarenes show only moderate affinity for the binding of smaller molecules and ions. A particularly interesting group of calixarene derivatives with proton donating groups on their substituents are secondary amide derivatives.^{1,2}

Cyclopeptides are macrocyclic compounds that contain amide groups within their backbone, and sometimes on their sidechains. Due to this, they are sometimes used as anion receptors.^{3,4} Their good complexation properties are a result of the proton-donor properties of peptide groups and the flexibility of the macrocyclic ring.

By combining the calixarene part, which determines the rigid conformation of the molecule and the cyclopeptide part, which carries the proton donor groups essential for the anion coordination, we created receptors that could have high affinity, and possibly selectivity, towards anions in the solution (Figure 1). Here we present the results of molecular dynamics simulations of several calixarene-cyclopeptide derivatives complexed with halogen and oxoanions in solution. We observed stable complexes in which the anions were coordinated by amide protons of calixarene and cyclopeptide ring.




Figure 1. Calixare-cyclopeptide receptor in complex with dihydrogen phosphate.

ACKNOWLEDGEMENTS This work was supported by the Croatian Science Foundation under project IP-2019-04-9560 (MacroSol). Scientific resources of Center of Excellence in Chemistry were used in this research.

REFERENCES

1. G. Horvat, V. Stilinović, B. Kaitner, L. Frkanec, V. Tomišić, *Inorg. Chem.* **2013**, *52*, 12702.
2. G. Horvat, L. Frkanec, N. Cindro, V. Tomišić, *Phys. Chem. Chem. Phys.* **2017**, *9*, 24316.
3. G. Horvat, S. Tarana, N. Vidović, N. Cindro, G. Speranza, V. Tomišić, *J. Mol. Liq.* **2021**, *340*, 116848.
4. I. Petters, M. Modrušan, N. Vidović, I. Crnolatac, N. Cindro, I. Piantanida, G. Speranza, G. Horvat, V. Tomišić, *Molecules* **2022**, *27*, 3918.

INFLUENCE OF DIFFERENTLY STABILIZED SILVER NANOPARTICLES ON CALCIUM PHOSPHATE FORMATION AND TRANSFORMATION

Suzana Inkret,^{a,*} Nadežda Radmilović,^b Kristina Smokrović,^c Nadica Maltar Strmečki,^c
Maja Dutour Sikirić^a

^a Laboratory for Biocolloids and Surface Chemistry, Division of Physical Chemistry, Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia, ^b Vinča Institute of Nuclear Sciences, 11001 Belgrade, Serbia ^c Laboratory for Electron Spin Spectroscopy, Division of Physical Chemistry, Ruđer Bošković Institute, Bijenička c. 54, 10000 Zagreb, Croatia

* sinkret@irb.hr

Calcium phosphate (CaP) composites with antimicrobial metal/metal oxide nanoparticles are emerging as a promising replacement for antibiotic therapy in implant-related infections. Silver nanoparticles (AgNPs) are attracting special attention due to their nonselective antibacterial activity.¹ A promising method of preparing CaP composites with AgNPs is precipitation.

The effect of differently stabilized silver nanoparticles on the formation and transformation of calcium phosphates was investigated. Polyvinylpyrrolidone (PVP), citrate (Cit) and sodium bis(2-ethylhexyl) sulfosuccinate (AOT) were used as stabilizers. Different effects on the rate of amorphous calcium phosphate (ACP) transformation to crystalline phase, as well as properties of formed amorphous and crystalline solid phases were observed depending on the type of AgNPs. After 60 min ageing time, in the control system ACP was transformed into mixture of calcium deficient hydroxyapatite (CaDHA) and small amount of octacalcium phosphate (OCP), as confirmed by PXRD, TEM, and SEM analysis of the formed precipitate. AOTAgNPs inhibited ACP transformation and OCP formation, at concentrations of 10 and 25 ppm. Furthermore, CitAgNPs at a concentration of 5 ppm promoted ACP transformation, but the effect on the composition of the obtained solid phase was not evidenced. No effect of PVPAgNPs on rate of transformation and precipitate composition was detected. The behaviour in the local environment was monitored by electron paramagnetic spectroscopy. SEM micrographs revealed that the best incorporation into CaPs was achieved for AOTAgNPs.

The obtained results provide new insights important for understanding CaPs formation on different nanomaterials.

ACKNOWLEDGEMENTS Financial support from the Croatian Science Foundation, Grant HRZZ- IP-2018-01-1493 is greatly acknowledged.

REFERENCES

1. M. Rai, A. Yadav, A. Gade, *Biotechnol. Adv.* **2009**, *27*, 76.

GLASS SURFACE MODIFICATION WITH INKJET PRINTABLE AMPHIPHILIC SILVER NANOPARTICLES

Irena Ivanišević,* Marko Zubak, Petar Kassal

Department of General and Inorganic Chemistry, Faculty of Chemical Engineering and Technology,
University of Zagreb, Marulićev trg 19, 10000 Zagreb, Croatia

* ivanisevic@fkit.hr

Additive manufacturing is one of the most rapidly adopted technologies today because it provides an efficient process for manufacturing devices. The ability to inkjet print thin films onto glass substrates is attractive for the application in electronics (i.e., printing active layers in sensors and memory devices), or in architecture and interior design to produce decorative glass surfaces.^{1,2} Inkjet printing is able to produce sophisticated and complex designs, as the process is fully digitally controlled and involves effective control at the micro-scale.

In this work, formulation and characterization of an ink containing amphiphilic silver nanoparticles (AgNPs) for the modification of glass surfaces by inkjet printing is presented. The AgNPs were prepared by modifying the primary particle stabilizer³ with functional amine (3-morpholinopropylamine) via an amidation reaction. The amphiphilic particle coating balances the hydrophilic-hydrophobic equilibrium and improves the dispersibility in less polar solvents, as well as adhesion of the printed features to clean glass surfaces in the fabrication of high-resolution metal patterns. Highly stable nanoink was prepared by redispersing 10 wt % AgNPs in a ternary solvent mixture of water, ethanol and ethylene glycol. The main characteristics of the formed ink, such as particle size and size distribution, stability over time and rheological properties were studied in detail. Multiple layers were printed on microscope slides, and post-printing treated both thermally and with intense pulsed light (IPL). In addition, electrical properties were evaluated using a four-point probe. Thus, new fabrication technologies (inkjet printing) and post-processing treatments (photonic annealing) combined with innovative materials (AgNPs as inorganic pigments) enabled the modification of glass surfaces with a thin metallic film that can be used in printed electronics or as customized decorative products.

ACKNOWLEDGEMENTS This work is funded by the European Regional and Development Fund for the Industrial Research and Innovation (IRI) Project “*Research and development of innovative glass processing techniques*” (KK.01.2.1.02.0059)

REFERENCES

1. R. Trattnig, G. Cattaneo, Y. Voronko, G. C. Eder, D. Moor, F. Jamschek, T. Buchsteiner, *Sustainability* **2021**, *13*, 12775.
2. A. A. Simon, B. Badamchi, H. Subbaraman, Y. Sakaguchi, L. Jones, H. Kunold, I. J. van Rooyen, M. Mitkova, *Sci. Rep.* **2021**, *11*, 14311.
3. I. Ivanišević, S. Milardović, P. Kassal, M. Zlatar, *Electrochim. Acta* **2021**, *377*, 138126.

UTILIZATION OF A KINETIC ISOTOPE EFFECT TO DECREASE DECOMPOSITION OF CEFTRIAZONE IN A MIXTURE OF D₂O/H₂O

Ivona Jasprica,^{a,*} Petar Horvat,^a Katarina Zrnc,^a Karl Bonney,^b Vidar Bjornstad,^b
Ernest Meštrović,^a Robert Vianello,^c Lucija Hok,^c Nikola Bregović,^d Josip Požar,^d Katarina Leko,^d
Vladislav Tomišić^d

^aXellia Ltd., Slavonska avenija 24/6, Zagreb, Croatia, ^bXellia Pharmaceuticals AS, Silurveien 2,
0380 Oslo Norway, ^cLaboratory for the Computational Design and Synthesis of Functional Materials,
Ruđer Bošković Institute, Bijenička c. 54, Zagreb, Croatia, ^dDepartment of Chemistry, Faculty of Science,
University of Zagreb, Horvatovac 102a, Zagreb, Croatia

* Ivona.jasprica@xellia.com

The labile β -lactam ring of penicillins and other β -lactam antibiotics is characterized by its pronounced susceptibility to various nucleophiles, acid-base reagents, oxidizing agents or even solvents like water and alcohol. However, the major pathways of β -lactams degradation are similar, leading to various breakdown products. The discovery of cephalosporin and demonstration of its improved stability in aqueous solution, as well as its enhanced *in vitro* activity against penicillin-resistant organisms, were major breakthroughs in the development of β -lactam antibiotics. Although cephalosporins are more stable with respect to hydrolytic degradation than penicillins, they still experience a variety of chemical and enzymatic transformations.

The present study was designed to gain insight into the kinetics of cephalosporin degradation, more specifically, ceftriazone degradation at its therapeutic concentration. As ceftriazone is one of the most frequently used parenteral antibiotics in treating specific infections, understanding of its degradation mechanisms at the concentration administered parenterally is essential for development of formulations containing this API.

Therefore, our study was focused on obtaining information on the rates and mechanisms of ceftriazone degradation in aqueous environment at the therapeutic concentration of 20 mg/mL. The study was directed towards obtaining kinetic information on the rates and mechanisms of ceftriazone degradation in water, a mixture of water and deuterium oxide, and deuterium oxide itself at the neutral pH (range of 6.7 to 7.6). Specific ceftriazone degradation products were observed in aged samples prepared in the examined solvents using HPLC and MS. By comparing the degradation rates in H₂O and D₂O, the observation of a kinetic isotope effect (KIE) provided some valuable insight as to the nature of the initial ceftriazone degradation. This result, in combination with an investigation of the reaction order for the degradation using the method of initial rates, highlighted the important contribution of the formation of a previously unreported dimer-type species. Computational analysis was utilized to get a molecular insight into chemical processes governing the ceftriazone degradation and to rationalize the stabilizing effect of replacing H₂O with D₂O. For that purpose, molecular dynamics simulations in both explicit solvents were carried out, together with a range of mechanistic DFT calculations to obtain the underlying kinetic and thermodynamic parameters of the most prevailing reactions in solution. In doing so, focus was on two dominant processes, (i) the opening of the β -lactam ring following the hydrolytic cleavage of its C–N amide bond, and (ii) the breaking of the C–S bond linking the triazine moiety with the rest of the structure.

ACKNOWLEDGEMENTS: The results presented here are co-financed by the European Union from the European Regional Development Fund (Liquid antibiotics and active substances with antibiotic-like stabilization mechanisms for intravenous and intramuscular application, KK.01.2.1.02.0175).

GLUTEN IN DIETARY SUPPLEMENTS USED IN THE TREATMENT OF INFLAMMATORY BOWEL DISEASES

Ilija Klarić,^a Jelena Kovačić,^b Mario-Livio Jeličić,^{b,*} Ana Mornar,^b Snježana Zubčić,^c
Daniela Amidžić Klarić^b

^a Department of Health Ecology, Public Health Brčko DC, R. Dž. Čauševića 1, Brčko DC, Bosnia and Herzegovina, ^b Department of Pharmaceutical Analyses, Faculty of Pharmacy and Biochemistry, A. Kovačića 1, Zagreb, Croatia, ^c Official Medicines Control Laboratory Division, Agency for Medical Products and Medical Devices of Croatia, Ksaverska cesta 4, Zagreb, Croatia

* mljelicic@pharma.hr

Gluten is naturally found in food, especially in certain grains such as wheat, rye, and barley. The consumption of gluten causes damage to the lining of the small intestine, preventing the absorption of nutrients and causing deficiencies that can lead to severe conditions and diseases. Inflammatory bowel diseases, Crohn's disease and ulcerative colitis, are chronic relapsing diseases affecting millions worldwide and disrupting their daily lives and requiring prescribing of drugs and dietary supplements. In addition to the active substance, orally formulated drugs contain a whole range of ingredients, such as excipients. Excipients can be synthetic or from natural sources originating from plants. One common excipient is starch. Corn, potatoes, and wheat are the most common starch types. Therefore, oral drugs and dietary supplements in the form of tablets and capsules are potential sources of gluten contamination. Plants and dietary supplements are important sources of biologically active components that have a positive effect on human health. In addition, the use of dietary supplements is constantly increasing. On the other hand, health professionals do not recognize the potential sources of gluten in orally formulated drugs. For all the listed reasons, this work aimed to evaluate gluten content in dietary supplements.

Forty-five samples were collected from local pharmacies, food health stores and purchased from the Internet (products online-only available in Croatia). All dietary supplements were classified as botanical monopreparations and multipreparation, as they contained one or more of the investigated herbal extracts: Indian frankincense, Green chiretta and Turmeric. Enzyme-linked immunosorbent assay (ELISA) was used to detect and quantitate gluten in investigated samples. All samples were analyzed in duplicate prior to stated expiry date.

Based on the obtained results, the gluten content was above the LOQ value (0.15 ng/mL) in 12 samples, while in one dietary supplement was over 20 ppm (28 µg/mL). Therefore, there are currently no labeling requirements for gluten found in orally formulated drugs and dietary supplements, while all foods labeled "gluten-free" must contain less than 20 ppm gluten.

To sum up, all analyzed dietary supplements except one sample contained gluten less than the FDA criteria.

ACKNOWLEDGEMENTS This work has been supported in part by the Croatian Science Foundation under the project numbers [UIP-2017-05-3949, DOK-2021-02-7922].

OPTIMIZATION OF LC-QqQ-MS/MS METHOD FOR QUANTIFICATION OF ANDROGRAPHOLIDES IN HUMAN PLASMA

Jelena Kovačić, Mario-Livio Jeličić, Daniela Amidžić-Klarić, Ana Mornar

Department of Pharmaceutical Analyses, Faculty of Pharmacy and Biochemistry, A. Kovačića 1, Zagreb

* mljelicic@pharma.hr

Andrographolides are diterpenoids found in the plant *Andrographis paniculata* and are widely used in traditional medicine, exhibiting anti-inflammatory properties.¹ In this research we developed and validated high-performance liquid chromatography coupled with electrospray ionization triple-quadrupole tandem mass spectrometry (LC-QqQ-MS/MS) method for quantification of andrographolides in human plasma. The chromatographic analysis of pharmacological active andrographolide (AND), neoandrographolide (NAND) and 14-deoxy-11,12-didehydroandrographolide (DAND) was performed on a 15 cm HSS Cyano column with a gradient mobile phase of ultra-pure water and acetonitrile with the addition of formic acid (0.1%) as a modifier at a flow rate of 0.8 mL/min in 21 min.

The validation results showed acceptable linearity of the method ($r > 0.999$) for all analytes within the calibration range 0.7–10 µg/mL. The acceptable accuracy of the method was indicated by recovery of 90–107 %. The limit of quantification was found to be between 2 and 70 ng/mL while the limit of detection was found between 0.6 and 20 ng/mL. Multiple reaction monitoring (MRM) mode was optimized to quantify data under monitoring precursor-product ion transitions of m/z 351.1 → 105.0, 351.5 → 91.0, 351.1 → 79.0 and 351.1 → 77.0 for AND, m/z 481.6 → 319.0, 481.6 → 301.2, 481.6 → 289.2, 481.6 → 95.1 for NAND and m/z 333.1 → 314.9, 333.1 → 297.2, 333.1 → 257.3, 333.1 → 91.0 for DAND (Figure 1). This advanced method is simple, selective, accurate and cost-effective. As a result, this assay can be easily transferred in clinical settings for pharmacokinetic studies and therapeutic drug monitoring programs.




Figure 1. Total ion scan (TIC) of AND, NAND and DAND collected in dynamic MRM mode.

ACKNOWLEDGEMENTS This work has been supported in part by the Croatian Science Foundation under the project numbers [UIP-2017-05-3949, DOK-2021-02-7922] and project FarmInova (KK.01.1.1.02.0021) funded by the European Regional Development Fund.

REFERENCES

1. F. Algieri, A. Rodriguez-Nogales, M. E. Rodriguez-Cabezas, S. Risco, M. A. Ocete, J. Galvez, *Mediators Inflammation*, 2015, 179616.

ALL-SOLID-STATE FLEXIBLE AMMONIUM-SELECTIVE ELECTRODE BASED ON INKJET PRINTED GRAPHENE SOLID CONTACT

Željka Boček, Sara Krivačić,* Petar Kassal

Faculty of Chemical Engineering and Technology, University of Zagreb, Marulićev trg 19,
 10000 Zagreb, Croatia

* skrivacic@fkit.hr

Continuous chemical monitoring presents an imperative for environmental and healthcare-related analysis, often inaccessible to classical analytical techniques, such as HPLC and UV/Vis spectroscopy. Within the scope of this work, miniaturized planar electrochemical sensor platforms set up in an all-solid-state manner have been developed by inkjet printing. We developed, characterized and optimized the printing process of a planar all-solid-state ammonium ion-selective electrode (NH₄-ASS-ISE) (Figure 1). Silver connection pads were made by inkjet printing commercial Ag-ink on flexible polymeric substrates. For the solid contacts, we have previously formulated an inkjet printable water-ethanol-ethylene glycol graphene suspension.¹ The number of graphene overprints and the volume of the drop-cast ammonium-selective membrane were optimized with the aid of electrochemical impedance analysis, cyclic voltammetry and potentiometry. NH₄-ASS-ISE prepared by 50 graphene ink overprints and drop-cast with 120 μL of the ammonium-selective membrane demonstrated the best properties, in terms of sensitivity (64.3 mV/dec in the linear range from 10⁻⁴ M to 10⁻¹ M NH₄Cl), potential stability, reproducibility of measurements (n = 3), reversibility, and linearity (R² = 0.9980).




Figure 1. Photograph and reversibility measurements of the inkjet printed electrode with 50 overprints of graphene solid contact and 120 μL of the ammonium-selective membrane.

ACKNOWLEDGEMENTS This work was funded by the Croatian Science Foundation under grant UIP-2020-02-9139.

REFERENCES

1. M. Kralj, S. Krivačić, I. Ivanišević, M. Zubak, A. Supina, M. Marciuš, I. Halasz, P. Kassal, *Nanomaterials* **2022**, *12*, 2936.


DETERMINATION OF THE OPTIMAL CONDITIONS FOR THE BIODEGRADATION OF POLYVINYL CHLORIDE BY THE YEAST *CANDIDA PARAPSILOSIS* USING A FULL FACTORIAL PLAN

Marija Kuštro,* Dubravka Tavra, Kristina Bule, Viktorija Prevarić, Matija Cvetnić, Šime Ukić, Tomislav Bolanča, Martina Miloloža, Marinko Markić, Lidija Furač, Dajana Kučić Grgić

University of Zagreb, Faculty of Chemical Engineering and Technology, Marulićev trg 19,
10000 Zagreb, Croatia

* mkustro@fkit.hr

The introduction of plastic waste into the components of the environment is increasing. We find plastic particles smaller than 5 mm in the environment, which we call microplastics (MP). MP has been attracting increasing attention for years and is considered a new pollutant in the environment. During the production of plastic products, various additives are added to improve their chemical and physical properties.¹ As a new pollutant, MP needs to be removed from the environment, so recently research has been conducted to find possible solutions for its removal. A combination of physical, chemical and biological processes can enhance the removal of MP from the environment. Microorganisms have a good potential for MP biodegradation which is an environmentally acceptable process.² In this research, the optimal conditions for biodegradation were investigated according to the design of a full-factorial plan. For this purpose, the study was conducted at three different optical densities (OD) (0.1, 0.5 and 1.0), pH values (3, 5 and 7) and revolutions per minute (120, 160 and 200). During the 30 days of the biodegradation experiment, changes in OD, pH values and the total number of grown yeast colonies (CFU) were monitored. For a better insight into the biodegradation process, TOC analysis, HPLC/MS analysis of filtrate, and FTIR-ATR analysis of MP were carried out. The toxicity of the filtrate was determined using the bacterium *Vibrio fischeri*. The experimental results were processed by statistical analysis of variance (ANOVA) using a full factorial experimental design. This study showed that the optimal conditions for PVC biodegradation are the rpm, at low values (120 rpm), at a higher OD (1.0) and an intermediate pH (5).



ACKNOWLEDGEMENTS This work was done as part of the project "Application of advanced water treatment technologies for the removal of microplastics" (IP-2019-04-9661) funded by the Croatian Science Foundation.

REFERENCES

1. S. Čačko, E. Pančić, I. Zokić, M. Miloloža, D. Kučić Grgić, *Kem. Ind.* **2022**, *71*, 49.
2. M. Dusinska, E. Rundén-Pran, J. Schnekenburger, J. Kanno, *Adverse Eff. Eng. Nanomater.* **2017**, *3*, 51.

POTENTIOMETRIC SENSORS AND WATER SOLUTIONS – SOLUTION FOR SURFACTANT ANALYSIS

Dubravka Madunić-Čačić,^{a,*} Blaženka Čiško-Anić,^a Nikola Sakač,^b Marija Jozanović^c

^a Saponia Chemical, Pharmaceutical and Foodstuff Industry, Inc., 31000 Osijek, Croatia,

^b Faculty of Geotechnical Engineering, University of Zagreb, 42000 Varaždin, Croatia,

^c Department of Chemistry, University of Osijek, 31000 Osijek, Croatia

* dubravka.madunic-cacic@saponia.hr

Green analytical methods¹ for anionic, cationic and non-ionic surfactants determination are presented. In all determination home-made liquid membrane potentiometric sensors as indicator electrodes and potentiometric titrator of high sensitivity and precision were used. Potentiometric titration were performed in an aqueous sample solution^{2,3} (e.g., detergents, washing and cleaning agents, etc.) or directly (wastewater samples⁴) without any pretreatment.

Due to the flexibility of this analytical technique, it provides the possibility of adjusting the determination conditions (e.g., adjusting the pH value of the analytical solution) with the aim of removing the influence of interferents (e.g., polycarboxylates, betaines, amine-oxides, soaps, etc.) on the accuracy of the determination. Furthermore, these methods have a whole series of other properties that make them interesting for expanding the field of their application. Besides the fact that these methods belong to green methods, and enable the quantitative determination of surfactants, they also provide monitoring of their behavior in complex matrices that are very common in reality.

Generally, these electroanalytical methods are green methods for determination of anionic, cationic and non-ionic surfactants in real samples³ and waste waters,⁴ and alternative for standard methods used. Despite the fact that standard methods^{5,6} use harmful organic solvents (e.g., chloroform) and are environmentally unacceptable as a green method, they are still the most prevalent in practice today.

ACKNOWLEDGEMENTS Investigations were conducted in Saponia (Osijek, Croatia), in cooperation with partners from research institutions.

REFERENCES

1. P. Anastas, N. Eghbali, *Chem. Soc. Rev.* **2010**, 39, 301; A. Gałuszka, Z. Migaszewski, J. Namieśnik, *Trends Anal. Chem.* **2013**, 50, 78.
2. D. Madunić-Čačić, M. Sak-Bosnar, O. Galović, N. Sakač, R. Matešić-Puač, *Talanta* **2008**, 76, 259.
3. D. Madunić, M. Sak-Bosnar, R. Matešić-Puač, *Int. J. Electrochem. Sci.* **2011**, 6, 240.
4. M. Sak-Bosnar, D. Madunić-Čačić, Z. Grabarić, B. Grabarić, *The Handbook of Environmental Chemistry*, Springer **2015**, 31, 157.
5. **ISO 2271:1989** Surface active agents - Detergents - Determination of anionic-active matter by manual or mechanical direct two-phase titration procedure; **ISO 2871-1:2010** Surface active agents - Detergents - Determination of cationic-active matter content - Part 1: High-molecular-mass cationic-active matter.
6. **ISO 7875-1:1996** Water quality - Determination of surfactants - Part 1: Determination of anionic surfactants by measurement of the methylene blue index (MBAS); **ISO 7875-2:1984** Water quality - Determination of surfactants - Part 2: Determination of non-ionic surfactants using Dragendorff reagent.

SYNTHESIS AND SOLVATOCHROMISM OF THREE NOVEL BROOKER'S MERCOCYANINE ANALOGUES

Antonio Magnabosco,^{a,*} Gordana Majstorović,^b Tomislav Portada^c


^a Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, HR-10000 Zagreb, Croatia, ^b Science School Vladimir Prelog, Vukovarska 269, HR-10000 Zagreb, Croatia, ^c Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička c. 54, HR-10000 Zagreb, Croatia

* amagnabosco@chem.pmf.hr

Merocyanine dyes are well known for its solvatochromism. In early 20th century the suggestion was given that merocyanine dyes can be used as solvent property indicators. A typical merocyanine, known as Brooker's merocyanine or systematically 1-methyl-4-[oxocyclohexadienylidene]ethylidene]-1,4-dihydropyridine (**MOED**), exhibits pronounced color changes with changes of solvent polarity: it goes from yellow, through red and violet, to blue in water, ethanol, acetone, and chloroform, respectively.¹ In this work we synthesized a few novel analogues of **MOED** by making slight changes in molecular structure. Three novel merocyanine dyes were synthesized by introducing structural changes on both aromatic rings of **MOED** molecule. All of synthesized compounds shows solvatochromic properties with great differences in color (Figure 1). By dissolving equal amount of four merocyanines, **MOED** and three novel ones, different color shifts occur, **MOED** and **AM-02** are showing hipsochromic (blue) shift and **AM-01** and **AM-03** bathochromic (red) shift. Synthesis of few additional **MOED** analogues, as well as their structural and spectroscopic studies are in progress.



Figure 1. Brooker's merocyanine in different solvents.



REFERENCES

1. L. G. S. Brooker, A. C. Craig, D. W. Heseltine, P. W. Jenkins, L. L. Lincoln, *J. Am. Chem. Soc.* 1965, 87, 2443.


TEMPERATURE EFFECT ON COORDINATIVE, HALOGEN AND TETREL BONDS IN CRYSTAL STRUCTURE OBTAINED FROM LEAD THIOCYANATE AND 3-IODOPYRIDINE

Lucija Marić, Nikola Bedeković,* Vladimir Stilinović

Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, 10000 Zagreb, Croatia

* nbedekovic@chem.pmf.hr

The tetrel bond is a non-covalent interaction between Lewis base and a group 14 element acting as Lewis acid. As the halogen bond, interaction is formed between the σ -hole (electron-deficient region of the donor atom) and the electron rich functional group which is a part of the same or the other molecular entity. Crystal structures in which the same types of halogen, tetrel and coordinative bonds are simultaneously present are extremely rare, which has so far reduced the possibility of direct comparison of those interactions. In this work, we have prepared single crystals of $\{[\text{Pb}_2(\text{SCN})_4(\mathbf{3Ipy})_4](\mathbf{3Ipy})_2\} \cdot \{[\text{Pb}(\text{SCN})_2(\mathbf{3Ipy})_2](\mathbf{3Ipy})\} \cdot [\text{Pb}_2(\text{SCN})_4(\mathbf{3Ipy})_6]$ by slow cooling of lead thiocyanate and 3-iodopyridine solution in ethanol-water mixture. Asymmetric unit of the determined crystal structure consist of three different complexes in which lead atoms are hemi-coordinated with the corresponding coordination numbers of 4, 5 and 6. Regardless of the coordination number, lead atoms serve as tetrel bond donors toward either thiocyanate S or N atoms, as well as to the pyridine nitrogen of **3Ipy** fragment. Furthermore, iodine atoms (as good and proven halogen bond donors) participate in several types of halogen bonds with thiocyanate groups. Given the fact that coordination, halogen and tetrel bonding patterns have found in the same crystal structure, this compound has served as an excellent model system for studying the effect of temperature on the lengths and geometries of those types of interactions. Crystal structures of the model compound were determined at 14 different temperatures in the range between 170 and 300 K and it was found to be stable at any temperature of interest (no decomposition or polymorphic transition has noticed during the experimental procedure). Furthermore, by changing the temperature, various interaction length and geometry shifts were observed, which were mutually correlated and discussed.¹



ACKNOWLEDGEMENTS This research was supported by the Croatian Science Foundation under the project IP-2019-04-1868.

REFERENCES

1. G. Mahmoudi, S. K. Seth, A. Bauzá, F. I. Zubkov, A. v. Gurbanov, J. White, V. Stilinović, T. Doert, A. Frontera, *CrystEngComm* 2018, 20, 2812.

IMPURITIES IN LATANOPROST EYE DROPS: ANALYTICAL PROCEDURE LIFECYCLE MANAGEMENT AND QbD APPROACH IN THAT PROCESS

Marina Mavrinac,* Vlasta Bertone

Jadran Galenski Laboratorij, JGL d.d., Svilno 20 b, Rijeka, Croatia

* marina.mavrinac@jglpharma.com

Latanoprost (isopropyl (Z)-7-[(1R,2R,3R,5S)-3,5-dihydroxy-2-[(3R)-3-hydroxy-5-phenylpentyl]cyclopentyl]hept-5-enoate) is a drug molecule used to treat ocular hypertension and open angle glaucoma. It possesses a carbon-carbon double bond. Due to possible different orientations of atoms around that bond, latanoprost can be present in two isomeric forms: mainly, it is present as a *cis*-isomer, which is the active form; but under some circumstances, *trans*-isomer can also be present. This *trans*-isomer (isopropyl (5E)-7-[(1R,2R,3R,5S)-3,5-dihydroxy-2-[(3R)-3-hydroxy-5-phenylpentyl]cyclopentyl]-5-heptenoate), also known as Impurity F listed in European Pharmacopoeia (Ph. Eur.), is considered one of the latanoprost impurities that arise from the synthesis pathway.¹ Main degradation product of latanoprost is latanoprost acid ((Z)-7-[(1R,2R,3R,5S)-3,5-dihydroxy-2-[(3R)-3-hydroxy-5-phenylpentyl]cyclopentyl]hept-5-enoic acid), also known as Ph. Eur. Impurity H. This impurity is formed in solutions, but is actually a biologically active form of latanoprost.² To control the amount of impurities F and H present in the final drug product, a robust and stability indicating analytical method had to be developed.

Method development is the first phase in analytical method lifecycle, followed by method validation (phase 2), routine analysis (phase 3) and change management (phase 4). All four phases constitute a cyclic process, so any phase can be re-evaluated at any time if necessary.

For phase 1, the goal was to develop one method for analysis of both impurities (F and H), having in mind pKa and logD of analytes. The method was successfully developed and validated and has been used in JGL for routine quality testing. In other words, phases 1, 2 and 3 were successfully completed.

As part of the continuous analytical procedure lifecycle management, phase 1 was challenged once again by using a QbD (Quality by design) approach and testing the robustness of the method. Method robustness was already tested and confirmed during validation, but a traditional approach was used: changing one parameter at a time. By using a QbD approach, robustness was tested once again by changing multiple parameters simultaneously. After QbD driven experiments, a MODR (Method Operable Design Region) was defined, which provided a better knowledge on method performance and confirmed that the method is still suitable for its intended use.

REFERENCES

1. European Pharmacopoeia 10.8, Vol 2. Strasbourg, Council of Europe; 2022. Latanoprost, p. 2230.
2. B. Sjöquist, J. Stjernschantz, *Surv. Ophthalmol.* **2022**, *47*, S6.

DIVALENT CATION EFFECTS ON POLY(DIALLYLDIMETHYLAMMONIUM CHLORIDE)/POLY(SODIUM 4-STYRENSULFONATE) NANOFILM DEPOSITION

Mia Mesić,* Tin Klačić, Davor Kovačević

Department of Chemistry, Division of Physical Chemistry, Faculty of Science, University of Zagreb,
Horvatovac 102a, 10000 Zagreb, Croatia

* mmesic@chem.pmf.hr

Polyelectrolytes are macromolecules that consist of a repeating sequence of subunits that bear an electrolyte group. According to the charge they are divided into polycations, polyanions and polyampholytes, while according to the degree of dissociation in solution, they are divided into strong and weak. Polyelectrolyte multilayers are films that are formed by alternating adsorption of polycations and polyanions on a surface, most often using layer-by-layer method.¹ The properties of the deposited film depend on the experimental parameters such as used polyelectrolytes, type of substrate, ionic strength, and type of background salt.^{2,3} In this study, we investigated how the type of divalent cation affects the multilayer prepared from poly(diallyldimethylammonium chloride) and poly(sodium 4-styrenesulfonate). The examined cations were Mg^{2+} , Ca^{2+} , Sr^{2+} , Ba^{2+} , Ni^{2+} , Zn^{2+} and Cu^{2+} . Ellipsometry was used for monitoring the growth of the films up to ten layers, while atomic force microscopy was used to determine morphology and surface roughness. The change in surface wettability during polyelectrolyte adsorption was monitored by tensiometry.

While for alkaline earth metal cations ion-specific effect was observed, for transition metal cations this was not the case. The thickness, roughness, and wettability of films prepared in the presence of transition metal cations were not significantly different. In contrast, these properties differed for multilayers prepared in the presence of alkaline earth metal cations.

ACKNOWLEDGEMENTS This research was supported by the Croatian Science Foundation under the bilateral Slovenian-Croatian APPLPEMS project (IPS-2020-01-6126).

REFERENCES

1. G. Decher, J. D. Hong, J. Schmit, *Thin Solid Films* **1992**, 210/211, 831.
2. J. Borges, J. F. Mano, *Chem. Rev.* **2014**, 114, 883.
3. M. Mesić, T. Klačić, A. Abram, K. Bohinc, D. Kovačević, *Polymers* **2022**, 14, 2566.

CHARGE DENSITY OF TETRACYANOETHYLENE (TCNE) RADICAL ANION

Valentina Milašinović^{a,*} Krešimir Molčanov,^a Petar Štrbac^b

^a Division of Physical Chemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia,

^b Division of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia

* valentina.milasinovic@irb.hr

π -Stacking interactions of organic radicals and charge-transfer compounds are responsible for their unique properties and have recently been employed in design of optoelectronics, magnetic, and conductive molecular materials. Stacking of planar organic radicals involves some of the closest interplanar and intermolecular C–C contacts, of *ca.* 2.9 Å.¹ Recent quantum mechanical models proposed that unusually short and strong interactions between planar radicals have a partial covalent character, and this type of weak multicentre covalent bonding has been dubbed '*pancake bonding*'. The strongest interaction of this type is found in pancake bonded dimers of radicals, and its energy can exceed 15 kcal mol⁻¹.²

The simplest and also one of the most thoroughly studied case of pancake bonding is two-electron four-centre covalent bond between tetracyanoethylene (TCNE⁻) radicals.³ The π -bonded tetracyanoethylene dimer has been investigated for the first time by the experimental X-ray charge density and this study will thus serve as a standard for comparison with other types of multicentre bonding. A detailed X-ray charge density study of TCNE radicals with a short separation distance of 2.81 Å revealed two symmetry-independent bonding critical points (3, -1) between the TCNE⁻ radical rings with maximum electron density of 0.17 e Å⁻³, and one ring critical point (3, +1) (Figure 1).




Figure 1. Critical points in a stack of TCNE⁻ radical anions. Bond paths are shown as red lines, bond (3, -1) critical points are shown as red spheres, and ring (3, +1) critical points are green.

ACKNOWLEDGEMENTS

The work was financed by the Croatian Science Foundation, grant no. IP-2019-04-4674.

REFERENCES

1. K. Molčanov; B. Kojić-Prodić, *IUCrJ* **2019**, *6*, 156.
2. K. Molčanov, V. Milašinović, B. Kojić-Prodić, *Cryst. Growth Des.* **2019**, *19*, 5967.
3. Z.-H. Cui, H. Lischka, T. Mueller, F. Plasser, M. Kertesz, *ChemPhysChem* **2014**, *15*, 165.

THERMODYNAMICS OF ANION BINDING BY PENTAPHENYLALANINE IN ACETONITRILE

Matija Modrušan,^{a,*} Lucija Otmačić,^a Nikolina Vidović,^b Nikola Cindro,^a Ivo Crnolatac,^c
Giovanna Speranza,^d Gordan Horvat,^a Vladislav Tomišić^a

^a Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia,

^b Department of Agriculture and Nutrition, Institute of Agriculture and Tourism, Karla Huguesa 8, 52440 Poreč, Croatia, ^c Department of Organic Chemistry and Biochemistry, Ruđer Bošković Institute, Bijenička cesta 54, Zagreb, Croatia, ^d Department of Chemistry, University of Milan, Via Golgi 19, Italy

* mmodrusan@chem.pmf.hr

Cyclic peptides represent a large and underexploited class of drug discovery candidates. Their improved metabolic stability and binding affinity, in comparison with linear analogues, result in greater resistance to enzymatic hydrolysis, increased bioavailability, and bioactivity.¹ In the synthesis of cyclic peptides, the main factor that affects the yield of a ring-closure reaction is the conformational preorganization, i.e., the ability of a linear precursor to adopt quasi-cyclic conformation in which the N- and C- termini are in a close proximity. One approach to achieve that is to use templating agents, such as anions, that can bind to linear precursors as well as cyclic peptides.^{2,3} In the scope of this work, we studied the binding of inorganic anions to pentaphenylalanine methyl ester (Figure 1) in acetonitrile by means of ¹H NMR, fluorimetric, and CD titrations. In addition, we carried out the MD simulations to get insight into the structural changes which occur upon anion binding to pentapeptide.




Figure 1. Structure of linear pentaphenylalanine methyl ester.

ACKNOWLEDGEMENTS This work was supported by the Croatian Science Foundation under project IP-2019-04-9560 (Macrosol) and European Regional Development Fund (project CluK, KK.01.1.1.02.0016).

REFERENCES

1. E. H. M. Mohammed, S. Lohan, R. K. Tiwari, K. Parang, *Eur. J. Med. Chem.* **2022**, *235*, 114278.
2. N. Vidović, G. Horvat, D. Riva, T. Rinkovec, N. Cindro, V. Tomišić, G. Speranza, *Org. Lett.* **2020**, *22*, 2129.
3. I. Petters, M. Modrušan, N. Vidović, I. Crnolatac, N. Cindro, I. Piantanida, G. Speranza, G. Horvat, V. Tomišić, *Molecules* **2022**, *27*, 3918.

DIMERIC CAPSULE VS. COLUMNAR POLYMER: STRUCTURAL FACTORS DETERMINING THE SOLUTION AND SOLID-STATE AGGREGATION BEHAVIOR OF AMINO ACID FUNCTIONALIZED BTA DERIVATIVES

Filip Perlitius,^{a,b,*} Anna Walczak,^{a,b} Miroslava Čonková,^{a,b} Grzegorz Markiewicz,^b Jack Harrowfield,^c Artur R. Stefankiewicz^{a,b}

^a Faculty of Chemistry, Adam Mickiewicz University, Uniwersytetu Poznańskiego 8, 61-614 Poznań, Poland, ^b Center for Advanced Technologies, Adam Mickiewicz University, Uniwersytetu Poznańskiego 10, 61-614 Poznań, Poland, ^c Institut de Science et d'Ingénierie Supramoléculaires, Université de Strasbourg, 8 Allée Gaspard Monge, 67083 Strasbourg, France.

* filip.perlitius@amu.edu.pl

Building blocks based on benzene-1,3,5-tricarboxamide (BTA) have been widely investigated, due to their simple synthesis and countless possibilities of modification. Very well-studied and extremely important group of BTAs is amino ester decorated molecules.¹ The amino acid moiety as a structural motif has major impact on the supramolecular structure, due to encoded H-bonding pattern, which generally leads to helical structures.²

Herein we present the non-covalent assembly of four benzene-1,3,5-tricarboxamide (BTA) derivatives formed from glycine or L-valine esters (-Me, -iPr), in the solid state and in chloroform solution by means of several analytical methods (NMR, FT-IR, CD, single crystal and powder XRD, TGA). Two types of self-assembled structures were characterized: dimeric capsule formed by NH...O=C- (ester) and columnar assembly with characteristic -NH...O=C- (amide) bond alongside with monomers.

In-depth solid state study revealed the crucial role of C-terminus protecting group structure to supramolecular aggregation in solid state, which has not been thoroughly investigated yet. The isopropyl ester group prevents the molecule from assembly into columnar structure, instead dimeric assembly is observed. On the other hand, it has no effect in chloroform solutions, where amino acid chain determines the self-assembly. Our study points out the essential influence of substituent in both – the alpha atom of the amino ester and the ester protecting group.

ACKNOWLEDGEMENTS We thank the National Science Centre (SONATA BIS 2018/30/E/ST5/00032) for financial support.

REFERENCES

1. A. Desmarchelier, B. G. Alvarenga, X. Caumes, L. Dubreucq, C. Troufflard, M. Tessier, N. Vanthuyne, J. Ide, T. Maistriaux, D. Beljonne, P. Brocorens, R. Lazzaroni, M. Raynal, L. Bouteiller, *Soft Matter* **2016**, *12*, 7824.
2. P. J. Stals, M. M. Smulders, R. Martin-Rapun, A. R. Palmans, E. W. Meijer, *Chem. Eur. J.* **2009**, *15*, 2071.

A NEW SOLID-STATE PROMETHAZINE HYDROCHLORIDE-SELECTIVE SENSOR BASED ON FUNCTIONALIZED MWCNT

Mateja Peršić,* Mirela Samardžić, Andrea Dandić, Mateja Budetić

Department of Chemistry, Josip Juraj Strossmayer University of Osijek, Cara Hadrijana 8/A,
31000, Osijek, Croatia

* mpersic@kemija.unios.hr

Promethazine hydrochloride (PM) is a phenothiazine derivative widely used as antihistaminic, antiemetic, analgesic and anticholinergic drug.¹ Also, it has an anesthetic and sedative effect, and can be used as a therapeutic agent for treating various mental disorders.² Regardless of its benefits, it has adverse effects, such as endocrinal, cardiac and reproductive alterations.³ Considering that, its determination in pharmaceutical formulations and biological samples is important. Different analytical methods can be used for PM determination but they commonly have drawbacks such as expensive instrumentation, time consuming and complicated procedures, and use of toxic solvents. Due to their simplicity, ion-selective electrodes (ISEs) could be a good alternative. In order to develop ISEs with better properties, ISE membrane can be modified with multi-walled carbon nanotubes (MWCNT) considering their unique properties.⁴

The new solid-state PM-selective sensor, with liquid type of membrane, was developed using MWCNT functionalized with a sulfate group and PM ion as a sensor material. The new sensor was characterized using direct potentiometry. Response characteristics, dynamic response, influence of pH and selectivity were determined.

ACKNOWLEDGEMENTS This study is supported by the European Structural and Investment Funds grant for the Croatian National Scientific Center of Excellence for Personalized Health Care, Josip Juraj Strossmayer University of Osijek (grant #KK.01.1.1.01.0010).

REFERENCES

1. S. S. Badawy, S. A. El Seoud El Said, *Am. J. Anal. Chem.* **2013**, *4*, 258.
2. O. Saleh, A. El-Azzouny, H. Aboul-Enein, A. M. Badawy, *Indian J. Pharm. Sci.* **2009**, *35*, 19.
3. T. Alizadeh, M. Akhoundian, *Electrochim. Acta* **2010**, *55*, 3477.
4. T. Yin, W. Qin, *TrAC, Trends Anal. Chem.* **2013**, *51*, 79.

CHARACTERIZATION OF SULFONYLUREAS' ADME PROPERTIES USING BIOMIMETIC CHROMATOGRAPHY AND COMPUTATIONAL PROCEDURES

Martina Rajić,^{a,*} Mario-Livio Jeličić,^b Jelena Kovačić,^b Daniela Amidžić Klarić,^b
Nikola Cindro,^a Ana Mornar Turk^b

^a Department of Chemistry, Faculty of Science, University of Zagreb, Horvatovac 102a, Zagreb, Croatia, ^b
Department of Pharmaceutical Analysis, Faculty of Pharmacy and Biochemistry,
University of Zagreb, Ante Kovačića 1, Zagreb, Croatia

* rajic.martina1@gmail.com

Sulfonylureas are the oldest class of oral antidiabetic medication used to treat type 2 diabetes. There are two generations of sulfonylureas, and both are equally effective in lowering glucose concentration in the blood. Still, they differ in pharmacological properties such as absorption, metabolism, elimination, and dosing. Therefore, modern and advanced methods are being researched daily to determine the pharmacological properties of the new potential drugs. One of the latest methods is biomimetic chromatography which uses columns that mimic biological conditions and computer (*in silico*) methods that predict various ADME (*Adsorption, Distribution, Metabolism, Excretion*) and QSPR (*Quantitative Structure-Property Relationship*) parameters.

The present study was designed to gain insight into the ADME parameters such as lipophilicity, solubility, phospholipophilicity, protein plasma binding, oral bioavailability, distribution coefficient, gastrointestinal absorption, and compliance with Lipinski's "rule of five" of three sulfonylureas (*N*-methylcarbamoyl)-4-methylbenzenesulfoamide, (*N*-benzylcarbamoyl)-4-methylbenzenesulfonamide, and 1,1'-trimethylenebis[3-(*p*-tolylsulfonyl)-urea] by using HPLC biomimetic chromatography and various computer programs for prediction of stated properties.

More precisely, our study focused on determining the lipophilicity and phospholipophilicity of three sulfonylureas using C18 (octadecylsilane) chromatographic and Immobilized Artificial Membrane (IAM) columns. Chromatographic columns with human serum albumin (HSA) and α -1-acid glycoprotein (AGP) were used to characterize plasma proteins binding. Various free and commercially available programs were used to predict the ADME properties of stated compounds. For the parameters for which it was possible, the experimentally obtained values were correlated with the calculated ones.

The experimental results showed that all three sulfonylureas are lipophilic, interact with phospholipids, and bind to HSA and AGP plasma proteins. Furthermore, a high coefficient of determination ($R^2 > 0.9$) was obtained for most of the calculated parameters with experimentally determined parameters. Therefore, obtained calculated and experimental results would benefit further research of the analyzed sulfonylureas as potential drugs.

RUTHENIUM(III) CATALYZED OXIDATIVE DEGRADATION OF PARACETAMOL BY HEXACYANOFERRATE(III) IN SOLUTION MEDIA – A COMPUTATIONAL SCREENING AND MECHANISTIC PATHWAY

Riya Sailani,* Asha Rolaniya

Department of Chemistry, Centre of Advanced Study, University of Rajasthan, Jaipur,
Rajasthan-302004 India

* l.p_riya@yahoo.co.in

The ruthenium (III) catalyzed oxidation of paracetamol (PCM) by hexacyanoferrate (III) has been studied in an acidic aqueous medium. The order in paracetamol is fractional but first-order with oxidant concentration. The reaction exhibits first order kinetics to ruthenium (III) and is catalysed by hydrogen ions in the form of $k_{obs} = a + b[H^+]$. The ionic strength does not affect the rate of reaction but is maintained in the reaction. The various activation parameters have been evaluated by employing Arrhenius and Eyring equation. Quinone oxime as the oxidation product of paracetamol has been confirmed spectrally. The oxidation reaction of paracetamol has been studied in eight different solvents and the rate data correlate satisfactorily with Kamlet-Taft's solvatochromic parameters (α , β , π^*) suggest that the specific solute-solvent interactions play a major role in governing the reactivity and the observed solvent effects have been explained on the basis of solute-solvent complexation. Based on the kinetic results, a suitable mechanism has been proposed.

To further support our proposed mechanism, density functional theory (DFT) computations at B3LYP/6-311*G(d,p) and LANL2MB show that activation energy barriers predict the same reactivity trend as shown by the kinetics experiments.

ACKNOWLEDGEMENTS This work is supported in part by the University Grant Commission, New Delhi, India through JRF for financial support. Acknowledgment is also extended to the co-authors and Head of the Department of Chemistry, University of Rajasthan Jaipur (Rajasthan, India). The authors would like to dedicate this work to (late) Prof. P. D. Sharma, Department of Chemistry, University of Rajasthan, Jaipur, India.

REFERENCES

1. P. Sharma, R. Sailani, A. Meena, C. L. Khandelwal, *J. Chem. Res.* **2020**, *44*, 295.
2. R. Sailani, D. Pareek, N. K. Soni, C. L. Khandelwal, *Curr. Phy. Chem.* **2014**, *4*, 290.
3. M. D. Martino, A. Chiarugi, *Pain Ther.* **2015**, *4*, 149.
4. C. Warwick, *J. Royal Society for Public Health* **2008**, 128(6).
5. R. Sailani, M. Sharma, D. Pareek, C. L. Khandelwal, P. D. Sharma, *Reac. Kinet. Mech. Catal.* **2012**, *105*, 249.
6. G. G. Graham, K. F. Scott, *Am. J. Ther.* **2005**, *12*, 46.
7. G. W. Przybyta, K. A. Szychowski, Gminski, *J. Clin. Exp. Pharmacol. Physiol.* **2021**, *48*, 3.
8. J. D. Sawant, K. K. Patil, G. S. Gokavi, *Trans. Met. Chem.* **2019**, *44*, 153.
9. R. Negi, B. Jain, S. Singh, A. K. Singh, A. Asthana. *SN Appl. Sci.* **2019**, *1*, 1380.

CYCLIC CARBAMATE FORMATION THROUGH PALLADIUM-CATALYSED CO₂ ABSORPTION BY PROPARGYLIC AMINES

Filip Vranješević,^a Nikola Sakač,^{b,*} Marija Jozanović,^c Maria Kolymjadi Markovic,^d
Valerije Vrčak,^d Dean Marković^a

^a University of Rijeka, Department of Biotechnology, Radmile Matejčić 2, 51000 Rijeka, Croatia, ^b Faculty of Geotechnical Engineering, University of Zagreb, Hallerova 7, 42000 Varaždin, Croatia, ^c University of Osijek, Department of Chemistry, Cara Hadrijana 8/A, HR-31000 Osijek, Croatia, ^d University of Rijeka, Faculty of Physics, Radmile Matejčić 2, 51000 Rijeka, Croatia, ^e University of Zagreb, Faculty of Pharmacy and Biochemistry, Ante Kovačića 1, 10000 Zagreb, Croatia

* nikola.sakac@gfv.unizg.hr

The rise of atmospheric CO₂ presents the biggest long-term challenge for the global climate and great efforts are being made to develop efficient and cost-effective CO₂ capture and storage technologies. In organic synthesis, this has led to an increased interest in using CO₂ as a cheap and readily available C1 synthon. Building on previously published work in this area,¹ we have synthesized several propargylic amine substrates and tested their ability to form carbamates through CO₂ absorption. The carbamate forming reaction was coupled to C-C bond formation using a palladium catalyst and several allyl-, aryl- and vinyl- substrates. The thermodynamics of the CO₂-absorbing reactions have been modelled using the Gaussian 16 software to find the most favorable substrates. We have optimized the reaction conditions to produce allylated carbamates in intermediate yields, finding that copper (I) co-catalysts can significantly improve the reaction yield. We have also synthesized several substrates for an intramolecular version of this reaction which will be explored at a later date. The mechanism of this reaction has also been modelled *in silico*, with preliminary NMR studies indicating that an unstable intermediate is formed by a reaction between the propargylic alcohol and a palladium(II) catalyst.

ACKNOWLEDGEMENTS We would like to thank HRZZ (Croatian Science Foundation) for funding this research as part of the research project IP-2019-04-8846 (In-silico potpomognut dizajn supstrata za hvatanje CO₂ u katalitičkoj formaciji karbonata i karbamata). F. Vranješević is employed and funded as part of the Croatian Science Foundation project “Career Development of Young Researchers-Training of New PhDs”.

REFERENCES

1. P. García-Domínguez, L. Fehr, G. Rusconia, C. Nevado. *Chem. Sci.* **2016**, 7, 3914.

SELECTION OF THE DELIVERY SYSTEM FOR THE PRESERVATIVE FREE OPHTHALMIC PRODUCT

Vesna Saršon,* Josipa Šumonja

Jadran Galenski Laboratorij, Svilno 20, 51000 Rijeka, Croatia

* vesna.sarson@jglpharma.com

There are specific challenges within the development of the primary packaging system in ophthalmic product without preservatives. In these products, the delivery system takes over the microbiological protection of the drug product instead of preservatives in the formulation. This brings a certain complexity that needs to be addressed.

In preservative free (PF) packaging, during the use, the product passes through dropper's chambers consisting of several different materials, prompting potential interactions, specifically in this case adsorption of the active substance.

It is known from literature data¹ that prostaglandins tend to adsorb to silicone material. Within this study, the adsorption of the prostaglandin was evaluated by measurement of assay in collected drops applied through the delivery system. Two prototypes were evaluated: Prototype 1 containing silicone parts and Prototype 2 without silicone parts.




Figure 1. Results of API assay measurements during in use period

(Figure 1). Prototype 2 did not present any major signs of API adsorption and results were in line with the assay specification limits (95 – 105 %).

Based on the results of the study, it was concluded that the prototype containing silicone parts is not a suitable delivery device for the product in development containing prostaglandine, due to the unsatisfactory level of adsorption of API, as described in literature data. This was not the case for the prototype without silicone parts which was selected as a better option for further studies.

REFERENCES

1. Y.Le Basle, P. Chennell, V. Sautou, *Pharm. Technol. Hosp. Pharm.* **2017**, *2*, 181.

DESIGN, SYNTHESIS AND CHARACTERIZATION OF LINEAR AND CYCLIC PEPTIDES

Sabino Sepčić,* Patrizia Janković, Marko Babić, Daniela Kalafatović

Department of Biotechnology, University of Rijeka, Radmile Matejčić 2, 51000 Rijeka, Croatia

* sabino.sepcic@gmail.com

Enzymes are proteins that help speed up chemical reactions by creating favorable environments for the conversion of substrates into products. The complexity of amino acid interactions and the large combinatorial space of proteins suggests that the earliest enzymes were short peptides. The peptides could have built the amyloid structures that provided the frameworks to sustain their catalytic activities. Inspired by a catalytically active peptide from Rufo et al.,¹ Ac-IHIHIQI-Am (Ac = acetyl *N*-terminal group, Am = amide *C*-terminal group), we have synthesized and characterized two linear and two cyclic peptides. The peptides were synthesized using the standard Fmoc solid phase peptide synthesis (SPPS). These peptides were tested for their supramolecular structure forming abilities in the presence of Zn²⁺. The linear peptides, Ac-IHIHINI-Am and Ac-IHINIHI-Am were synthesized to test the effect of changing the glutamine (Q) to asparagine (N) and the changing of position of asparagine within the sequence on the self-assembly and catalytic properties of the peptide. The cyclic variations of the linear peptides, cyclo-(IHIHINIE-Am) and cyclo-(IHINIHIIE-Am) were prepared through the *N*-terminal to glutamic acid side chain cyclization. The glutamic acid with the protecting group O-2-PhiPr was selectively deprotected on resin, after which the deprotected carboxyl group was coupled with the free *N*-terminus. By using cyclization to form intra-connected peptides, we expect that the increased rigidity introduced by cyclization will improve the catalytic efficiency of short peptides. The synthesized peptides were analyzed with liquid chromatography and mass spectrometry techniques. The data proved that the octapeptide on-resin cyclization was successful. Hydrogen bonding patterns were analyzed using infrared spectroscopy. The linear peptides showed an amide I parallel β -sheet-like structure in the infrared spectrum in the presence of Zn²⁺. Computational simulation was performed with the linear peptides to visualize the histidine coordination of Zn²⁺. The catalytic activity of the peptides will be assessed in future work.

REFERENCES

1. C. M. Rufo, Y. S. Moroz, O. V. Moroz, J. Stöhr, T. A. Smith, X. Hu, W. F. DeGrado, I. V. Korendovych, *Nat. Chem.* **2014**, *6*, 303.

THE BIOSURF PROJECT: A SUSTAINABLE PRODUCTION OF BIO-BASED SURFACTANTS FROM RENEWABLE RESOURCES

Giuseppe Cappelletti,^a Eleonora Pargoletti,^a Marco Rabuffetti,^a Marina Simona Robescu,^b Sara Sangiorgio,^a Riccardo Semproli,^b Daniela Ubiali,^b Giovanna Speranza^{a,*}


^a Department of Chemistry, Università degli Studi di Milano, via Golgi 19, 20133 Milan, Italy,

^b Department of Drug Sciences, University of Pavia, Viale Taramelli 12, 27100 Pavia, Italy

* giovanna.speranza@unimi.it

The project BioSurf aims at developing a self-sustainable productive chain of bio-based surfactants (sugar fatty acid esters, SFAEs) by fully upgrading the main waste stream of dairy industry, namely cheese whey permeate. Sugar fatty acid esters (SFAEs) are non-ionic surfactants that are characterized by excellent surface and interfacial tension reduction capability, low toxicity, and biodegradability. These features make SFAEs extremely promising for industrial applications as emulsifiers in the cosmetic and food sectors.¹ Interestingly, SFAEs can be obtained from renewable resources by enzymatic and/or chemoenzymatic approaches, thus answering the need for evermore sustainable and circular chemistry.²

6-*O*-Palmitoyl-1-*O*-butyl glucosides and galactosides were enzymatically synthesized by reacting *n*-butyl glucosides and galactosides, respectively, with molten palmitic acid in an easily scalable solvent-free system. Conversion of glucose and galactose into alkyl glycosides before the esterification reaction played a key role to circumvent the striking different solubility of sugars and fatty acids. In addition, 6-*O*-palmitoyl-1-*O*- β -D-butyl galactopyranoside was obtained by a two-step fully enzymatic approach, i.e., a transglycosylation reaction catalyzed by covalently immobilized β -galactosidase from *Aspergillus oryzae*, followed by Novozym[®] 435-catalyzed esterification. The physico-chemical properties of these tensides, such as interfacial tension features, W/O emulsification capability and W/O stability over time were deeply investigated.^{3,4}



ACKNOWLEDGEMENTS This work was financially supported by Cariplo Foundation (Italy) (call: “Circular Economy for a sustainable future 2020”, project BioSurf, ID 2020-1094).

REFERENCES

1. N. S. Neta, J. A. Teixeira, L. R. Rodrigues, *Crit. Rev. Food Sci. Nutr.* **2015**, *5*, 595.
2. T. Keijer, *Nat. Chem.* **2019**, *11*, 190.
3. S. Sangiorgio, E. Pargoletti, M. Rabuffetti, M. S. Robescu, R. Semproli, D. Ubiali, G. Cappelletti, G. Speranza, *Colloid Interface Sci. Commun.* **2022**, *48*, 100630.
4. R. Semproli, M. S. Robescu, S. Sangiorgio, E. Pargoletti, T. Bavaro, M. Rabuffetti, G. Cappelletti, G. Speranza, D. Ubiali, *submitted*.

STACKING PATTERNS OF ORGANIC RADICALS IN TETRACYANOQUINODIMETHANE SALTS

Petra Stanić,* Krešimir Molčanov

Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia

* pstanic@irb.hr

Crystals of planar organic radicals are interesting because of pronounced π -stacking interactions. π -stacking of aromatic rings is usually weak interaction, however, interactions between planar radicals are much stronger.¹ Recent studies in crystal engineering show that energy of some π -contacts may be comparable to stronger intermolecular interactions such as hydrogen and halogen bonding. Close contact of two planar radicals usually involves spin coupling so the energy of such contacts may be as high as $-20 \text{ kcal mol}^{-1}$.² Furthermore, a long infinite stack of equidistant radicals contributes to special properties of such crystals like semiconductivity and antiferromagnetism.² Therefore, these compounds have an important role in crystal engineering of functional materials such as optoelectronics, magnetic, and conductive molecular materials.^{1,3}

Here we report a study on π -stacking and multicentre bonding in salts of 7,7,8,8-tetracyanoquinodimethane (TCNQ) radical anions with different organic cations: tetrazolium (**1**), 1,4-dimethyl-DABCO cation (**2**), *N,N*-dipyridylmethanium (**3**), *N,N*-dimethyl-4,4'-bipyridinium (**4**), benzyl-*N*-methyl-4-pyridyl quinone (**5**) and *N,N,N',N',N',N'*-hexamethyl ethylenediamine (**6**) cations. Prepared crystals were studied by means of X-ray crystallography. In studied compounds stacking interactions are predominant with interplanar separations between rings of 3,0 – 3,2 Å. TCNQ moieties form dimers, stacks of dimers, trimers or tetramers with alternating shorter and longer separation between TCNQ rings or stacks in columns with equidistant separations (Fig 1). Besides that, in some salts N–H \cdots N and C–H \cdots N hydrogen bonds are present.




Figure 1. Different types of stacking of TCNQ moieties in prepared salts: a) and b) dimers, c) trimers, d) tetramers and e) equidistant stack.

ACKNOWLEDGEMENTS The work was financed by the Croatian Science Foundation, grant no. IP-2019-04-4674.

REFERENCES

1. K. Molčanov, B. Kojić-Prodić, *IUCrJ* **2019**, *6*, 156.
2. V. Milašonović, A. Krawczuk, K. Molčanov, B. Kojić-Prodić, *Cryst. Growth Des.* **2020**, *20*, 5435.
3. K. Molčanov, V. Milašonović, B. Kojić-Prodić, *Cryst. Growth. Des.* **2019**, *19*, 5967.

OPTIMIZATION STUDIES OF SOLID PHASE EXTRACTION AND LC-MS ANALYSIS OF CDK4/6 INHIBITORS IN BREAST CANCER COMBINATION THERAPY IN PLASMA SAMPLES

Lu Turković,^{a,*} Zvonimir Mlinarić,^a Kristina Javorić,^b Juraj Vuić,^b Anamarija Skenderović,^c
Sandro Makarić,^c Vladimir Radić,^c Tajana Silovski,^d Dragana Mutavdžić Pavlović,^b
Miranda Sertić^a

^a Department of Pharmaceutical Analysis, University of Zagreb Faculty of Pharmacy and Biochemistry, Ante Kovačića 1, Zagreb, Croatia, ^b Department of Analytical Chemistry, University of Zagreb Faculty of Chemical Engineering and Technology, Marulićev trg 20, Zagreb, Croatia, ^c GxR&D Analytics Zagreb, Global R&D, Teva Pharmaceuticals, Prilaz baruna Filipovića 25, Zagreb, Croatia, ^d Department of Oncology, University Hospital Centre Zagreb, Kišpatićeva 12, Zagreb, Croatia

* lu.turkovic@pharma.unizg.hr

Palbociclib, ribociclib and abemaciclib are novel anticancer agents used for HR+, HER2- breast cancer treatment in combination with anastrozole, letrozole or fulvestrant. They are prone to inter-individual variabilities and concentration-dependent adverse events, therefore better treatment outcomes might be achievable with therapeutic drug monitoring (TDM). One of the main prerequisites to TDM is the existence of a reliable and, preferably, fast, simple, and ecologically acceptable bioanalytical method. Few bioanalytical methods have been published so far for the determination of all these six drugs in plasma,^{1,2} however using only protein precipitation for sample clean-up. In this work, a solid-phase extraction (SPE) procedure was optimized for the simultaneous extraction of all six drugs of interest from plasma. Several SPE phases were tested: octylsilyl (C8, 200 mg/3 mL, 500 mg/3 mL), octadecylsilyl (C18, 200 mg/3 mL), hydrophilic-lipophilic balance (HLB, 60 mg/3 mL), mixed-mode cation exchange (MCX, 30 mg/1 mL), and mixed-mode weak cation exchange (WCX, 60 mg/3 mL). Analytes were eluted with methanol, acetonitrile, 2 % formic acid in methanol or 5 % ammonia in methanol. The samples were analyzed with Agilent 1290 Infinity II ultra-high performance liquid chromatograph coupled to Agilent 6470 triple quadrupole mass spectrometer. Waters XBridge Phenyl column (150 × 4.6 mm, 2.5 μm) was used as the stationary phase at 35 °C. The mobile phase consisted of water and methanol with 0.1 % formic acid in gradient elution at a 0.6 mL/min flow rate. The optimal solution was found to be the C8 phase (200 mg/3 mL), eluted with 1500 μL of methanol, which yielded extraction recoveries above 90 % for all analytes. This method was validated in terms of intra and inter-day precision and accuracy, linearity and calibration in the clinically relevant ranges, selectivity, carry-over, and matrix effects. The validated method was then successfully applied for the analysis of samples from patients taking different combinations of the drugs of interest. The proposed method is simple, fast, sensitive, providing good sample clean-up and high extraction yields for all six analytes of interest. Its application on real patient plasma samples supports its suitability for the purposes of further TDM testing.

ACKNOWLEDGEMENTS This work has been fully supported by the Croatian Science Foundation, under the project number UIP-2019-04-8461 and DOK-2021-02-4595.

REFERENCES

1. Y. Sato, K. Shigeta, T. Hirasawa, T. Sato, J. Ogura, M. Maekawa, A. Ebata, Y. Hamanaka, H. Tada, T. Ishida, M. Kikuchi, N. Mano, *J. Chromatogr. B* 2021, 1173, 12265.
2. L. Turković, L. Bočkor, O. Ekpenyong, T. Silovski, M. Lovrić, S. Crnković, B. Nigović, M. Sertić, *Pharmaceuticals* 2022, 15, 614.

NOVEL TECHNIQUES FOR SECONDARY STRUCTURE DETERMINATION OF PEPTIDES IN DRUG DEVELOPMENT

Dorina Vidović,* Nada Košutić Hulita, Nikola Marjanović

Research and Development, Pliva Hrvatska d.o.o., Prilaz Baruna Filipovića 25, 10000 Zagreb, Croatia

* Dorina.Vidovic@pliva.com

Knowledge of the peptide secondary structure is essential for determining its properties. The importance of secondary structure determination is also in peptide classification, function and structural change prediction and it is an important information for the tertiary structure. In general, secondary structure is a term that refers to local folded structures that form within a peptide due to interactions between atoms of the backbone. The two most common types of secondary structure are the α -helix and the β pleated sheet.¹

Today, regulatory agencies require detailed determination of peptide secondary structure. This is the reason why it is important to examine the secondary structure in detail when developing a new drug product in the pharmaceutical industry. It is necessary to determine the similarity of the new product and the reference drug product (RDP) in case of generic drug product development. Some of analytical techniques used in pharmaceutical industry for secondary structure determination are Raman spectroscopy, FTIR analysis, Circular Dichroism (CD) spectroscopy and Nuclear magnetic resonance (NMR).² Raman spectroscopy and FTIR analysis are techniques that determine the secondary structure based on the obtained spectrum in the area of amide I and amide II groups (characteristic area of secondary structure). CD spectroscopy measures the difference in absorbance of right- and left-circularly polarized light. Based on far UV spectra and mathematical calculations it is feasible to determine exact secondary structure.

This paper presents the results of secondary structure determination during drug product development using peptide active pharmaceutical ingredient (API). The secondary structure was tested on three lots of the new drug and on three lots of reference drug product. Methods were developed and samples were analyzed by Raman spectroscopy, FTIR analysis and CD spectroscopy. Analysis of RDP samples revealed the clear dominance of α -helix structure. Additionally, the percentage of α -helix structure in the range 50 – 57 % was determined. From the results obtained from the analysis of a new drug samples, the dominance and the percentage of the α -helix structure in range 50 – 57 % were also determined. Overall, the same secondary structure of all analyzed samples can be concluded, as well as the comparability of the results of all three analyses.

REFERENCES

1. L. Skipper, *Encyclopedia of Analytical Science*, Second Edition, **2005**, 412.
2. K. Wang, D.W Sun, H. Pu, Q. Wei, *Trends Food Sci. Technol.* **2017**, 67, 207.

DEEP EUTECTIC SOLVENT SYSTEM BASED ON AMINO ACIDS DERIVATIVES

Sanja Vukoja,^a Sandra Siročić,^a Jelena Ranić,^a Tina Medančić,^a Diana Žgela,^a Lucija Hok,^b
Ivona Jasprica,^a Robert Vianello,^b Ernest Meštrović^a

^a Xellia Ltd., Slavonska avenija 24/6, Zagreb, Croatia, ^b Laboratory for the Computational Design and Synthesis of Functional Materials, Ruđer Bošković Institute, Bijenička c. 54, Zagreb, Croatia

* sanja.vukoja@xellia.com

The focus of work presented here was to study and describe the following:

- (1) The effect of proline series molecules (L-Proline, *N*-acetyl-L-proline, L-prolinamide, *N*-acetyl-L-prolinamide) on the stability of piperacillin and tazobactam considering experimental results and molecular modelling data. This is a very interesting family of molecules, since all of them are based on L-proline, yet significantly different when it comes to their overall charges and protonation states. L-Proline is neutral, but zwitterionic, *N*-acetyl-L-proline is anionic, L-prolinamide is cationic, and *N*-acetyl-L-prolinamide is neutral without any charged fragments.
- (2) Utilization of deep eutectic solvent systems (DES) for stabilization of piperacillin and tazobactam, where one of the components is *N*-acetyl-L-prolinamide.

Furthermore, preparation and stability testing of formulations in the form of a deep eutectic solvent that comprise piperacillin and/or tazobactam and at least one *N*-acetyl amino acid amide was conducted. Some of the evaluated excipients were *N*-acetyl-D-alaninamide, and as mentioned above, *N*-acetyl-L-prolinamide.

It was discovered that the stability of the formulation is more enhanced in the specific DES systems. More precisely, DES compositions provided improved stability over time (evaluated through specific impurities increase and purity drop) when stored refrigerated (2-8 °C), and at room temperature (25 °C) in comparison to liquid (aqueous) formulations that only contain piperacillin and/or tazobactam. Computational analysis utilized a combination of quantum-chemical calculations and molecular dynamics simulations to characterize relevant interactions in solutions in relation with the studied degradation mechanisms, and to help in interpreting the observed experimental trends.

ACKNOWLEDGEMENTS: The results presented here are co-financed by the European Union from the European Regional Development Fund (Liquid antibiotics and active substances with antibiotic-like stabilization mechanisms for intravenous and intramuscular application, KK.01.2.1.02.0175).

NUCLEATION KINETICS OF ASCORBIC ACID IN WATER AND WATER-ETHANOL SOLVENT


Jyoti Yadav,^{a,b,*} Thomas Kendall,^b Carmen Guguta,^b Swati A. Patel^a

^a Indian Institute of Technology, Ropar, India, ^b Technobis Crystallizations Systems, Alkmaar, Netherlands

* jyoti.yadav@technobis.com

L-Ascorbic acid, commonly known as vitamin C, is an essential compound with a wide range of applications in pharmaceutical and health care industry, such as multivitamins, drug combination, nutritional supplement, and antioxidants in personal care products. Crystallization is an essential separation and purification step for the production of ascorbic acid. Furthermore, the solvent employed for crystallization plays a fundamental role in determining the crystal size and shape.¹ Nucleation is the first step of crystallization which shapes the crystal appropriately, and hence, it becomes crucial to understand the nucleation kinetics to control and optimise the crystallization process.

The present work determines the nucleation kinetics of ascorbic acid in water and water-ethanol binary solvent mixture (composition of 0.4:0.6 mole fraction), by isothermal method (induction time). Induction time is measured more than 80 times at four different supersaturations (S) using *Technobis Crystallization Systems' Crystal16 V3* instrument.^{2,3} The



experimentally measured induction time data points determine the probability to form nucleus in that timeframe. Nucleation kinetics is defined by classical nucleation theory, which relates the probability of nucleation with the supersaturation, represented by the equation

$$J = AS \exp\left(\frac{-B}{\ln^2 S}\right)$$

where A and B are kinetic and thermodynamic factors, respectively. Addition of ethanol in water solvent reduces the nucleation kinetics, indicating slower nucleation rate in the water-ethanol binary system. Ethanol reduces the polarity of the binary solvent mixture and increases the energy barrier to nucleate.

Solvent	A [$\text{m}^{-3} \text{s}^{-1}$]	B
Water	2868.25	1.587
0.6 Ethanol+0.4 Water	450.47	1.506

REFERENCES

1. S. Hassan, F. Adam, M. A. Bakar, S. A. Mudalip, *J. Saudi Chem. Soc.* **2019**, *23*, 239.
2. S. Jiang, J. H. ter Horst, *Cryst. Growth Des.* **2011**, *11*, 256.
3. C. Brandel, J. H. ter Horst, *Faraday Discuss.* **2015**, *179*, 199.

PHOTOREDUCTION IN SOLUTIONS BY INTENSE PULSED LIGHT

Marko Zubak,* Sara Krivačić, Petar Kassal

Faculty of Chemical Engineering and Technology, University of Zagreb, Marulićev trg 19,
 10000 Zagreb, Croatia

* mzubak@fkit.hr

Intense Pulsed Light (IPL) is a short pulsed flash (down to 100 μ s) of a xenon lamp emitting mostly visible light (190 nm to 1100 nm), which is caused by the formation of an electric arc between high-voltage tungsten electrodes (1000 V – 3000 V). Compared to other reduction methods, IPL photoreduction does not require using reducing agents; it is energy and time-efficient,¹ thereby meeting the requirements of sustainable development. In this study, IPL photoreduction was evaluated in two different systems (Figure 1). The first system was the dispersion of graphene oxide (GO) in water, buffer (pH= 10), ethanol, and ethanol with poly(vinyl butyral) (PVB). The purpose of the experiment was to determine the influence of the dispersing agent on photoreduction and to define the optimal parameters of the IPL system that provide a high-quality product. Additionally, the dispersion stability of the product was evaluated. The degree of reduction was verified by UV/Vis spectrophotometry; the largest degree of reduction and most stable dispersion was found in the buffer (pH= 10).

The second system was a methanol solution of AgCl and NaCl in the PVB polymer matrix. IPL partial photoreduction was performed at mild IPL parameters. Silver nanoparticles were formed, as evidenced by the peak at 420 nm in the UV/Vis spectra; with the excess of AgCl and NaCl still present in the PVB matrix. The obtained polymer-inorganic salts hybrid was used as a reference membrane (RM) in a proof-of-concept experiment for the development of a planar all-solid-state reference electrode. The RM has shown a great capability of ensuring potential stability in Cl^- and NO_3^- containing solutions, as well as in a broad pH range.




Figure 1. Scheme of the reduction of silver chloride and graphene oxide by intense pulsed light.

ACKNOWLEDGEMENTS: This work was supported by the Croatian Science Foundation (projects UIP-2020-02-9139 and DOK-2021-02-2362).

REFERENCES

1. Y.-R. Jang, S.-J. Joo, J. H. Chu, H.-J. Uhm, J.-W. Park, C.-H. Ryu, M.-H. Yu, H.-S. Kim, *Int. J. of Precis. Eng. and Manuf.-Green Tech.*, **2021**, *8*, 16.

SYNTHESIS OF NEW CATIONIC AMPHIPHILES WITH ANTIMICROBIAL ACTIVITIES

Kristina Đambić,^a Katarina Lubina,^a Dora Tot,^a Nikola Sakač,^{b,*} Dubravka Madunić-Čačić,^c
Dean Marković,^d Bojan Šarkanj,^e Iva Pukleš,^f Nada Glumac,^g Marija Jozanović^a

^a Department of Chemistry, Josip Juraj Strossmayer University of Osijek, 31000 Osijek, Croatia

^b Faculty of Geotechnical Engineering, University of Zagreb, 42000 Varaždin, Croatia ^c Saponia Chemical, Pharmaceutical and Foodstuff Industry, Inc., 31000 Osijek, Croatia ^d Department of Biotechnology, University of Rijeka, 51000 Rijeka, Croatia ^e Department of Food Technology, University North, 48000

Koprivnica, Croatia ^f Doctoral School of Chemistry, University of Pécs, 7624 Pécs, Hungary

^g Međimurske Vode Ltd., 40000 Čakovec, Croatia

* nsakac@gfv.unizg.hr

Cationic surfactants, such as pyridinium, benzimidazolium and imidazolium salts, have excellent biocidal activities against broad spectrum of bacteria and fungi. A series of new cationic amphiphiles of imidazolium and benzimidazolium with antimicrobial activities for potential development of new disinfectants and sanitizers were synthesized. Imidazolium and benzimidazolium ionic liquids containing long apolar alkyl and polyalkyl chains and different anionic part were prepared in the form of quaternary salts. Using one-step procedure, salts with an alkyl/dialkyl N-substituents and Br/large inorganic anions counter anions were prepared in good yields. New imidazolium and benzimidazolium based ionic liquids were characterized by FT-IR, NMR and LC-MS. Their chemical and spectral properties were discussed. The dependence of the antimicrobial activity on number of chains, chain length and anion will be determined in near future.

Keywords: amphiphiles, antibactericides, N-alkylation, nitrogen heterocycles, quaternary ammonium salts

EVALUATION OF EMBRYOTOXICITY OF GRANULATAMIDE B AND ITS STRUCTURAL ANALOGUES USING ZEBRAFISH MODEL

Dario Matulja^a, Gabrijela Matijević^b, Sanja Babić^b, Maria Kolymjadi Marković^c, Marija Jozanović^d, Nikola Sakač^{e,*}, Rozelindra Čož-Rakovac^b, Sandra Kraljević Pavelić^f, Dean Marković^a

^a University of Rijeka, Department of Biotechnology, Radmile Matejčić 2, 51000 Rijeka, Croatia

^b Laboratory for Aquaculture Biotechnology, Division of Materials Chemistry, Ruđer Bošković Institute, Bijenička cesta 54, 10000 Zagreb, Croatia

^c Faculty of physics, University of Rijeka, Radmile Matejčić 2, 51000 Rijeka, Croatia

^d University of Osijek, Department of Chemistry, Cara Hadrijana 8/A, 31000 Osijek, Croatia

^e Faculty of Geotechnical Engineering, University of Zagreb, Hallerova 7, 42000 Varaždin, Croatia

^f University of Rijeka, Faculty of Health Studies, Viktora Cara Emina 5, 51000 Rijeka, Croatia

* nikola.sakac@gfv.unizg.hr

Marine natural alkaloid, granulamide B, belonging to the class of *N*-fatty acyl tryptamines, have been isolated from the coral species of the *Eunicella* genus.¹ Despite the known moderate activity against various cancer cell lines *in vitro*, little is known about its other biological activities. Therefore, within our project, *Bioprospecting of the Adriatic Sea*, we have synthesized granulamide B according to the procedure of Fürstner's Fe-catalysed synthesis², accompanied with several structural analogues formed by modification of the acyl moiety. A small compound library has been designed, containing a total of 13 compounds differing in the number of C-atoms, saturation degree and conjugation of double bonds. The aim of this study was to evaluate the effects of the synthesized compounds towards the formation and development of zebrafish *Danio rerio* embryos following previously described protocol.³ Among tested compounds in our study, 4 out of 13 harmed the survival and embryonic development of *D. rerio* within the tested concentration range. The highest toxicity in terms of zebrafish mortality was observed for compounds containing fatty acyl chains with conjugated double bonds including granulamide B (LC₅₀=44.34 μM). In addition, the most common developmental abnormalities observed during exposure to above mentioned compounds were edemas, scoliosis and blood accumulation.

ACKNOWLEDGEMENTS We would like to thank Croatian Government and the European Union (European Regional Development Fund—the Competitiveness and Cohesion Operational Program - KK.01.1.1.01) for funding this research through project *Bioprospecting of the Adriatic Sea* (KK.01.1.1.01.0002) granted to The Scientific Centre of Excellence for Marine Bioprospecting - BioProCro. We also acknowledge the project “Research Infrastructure for Campus-based Laboratories at University of Rijeka”, co-financed by European Regional Development Fund (ERDF) and the University of Rijeka research grants UNIRI-prirod-18-102 and UNIRI-biomed-18-133. We would like to thank Croatian Science Foundation project ‘Career Development of Young Researchers-Training of New PhDs’ for funding PhD of D. Matulja and research project IP-2019-04-8846.

REFERENCES

1. Reyes F, Martín R, Fernández R. *Journal of Natural Products* 2006, 69, 4.
2. Sun CL, Fürstner A. *Angewandte Chemie* 2013, 52, 493. Babić S, Čižmek L, Maršavelski A. *Scientific Reports* 2021, 11, 1

AUTHORS

Adamska, Wiktorija	79	Gruić Sovulj, Ita	51
Alešković, Marija	80	Guguta, Carmen	120
Alić Stolar, Jasna	81	Hadži, San	28
Amidžić-Klarić, Daniela	98	Harrowfield, Jack	54, 108
Babić, Marko	114	Hok, Lucija	96, 119
Babić, Sanja	123	Horvat, Gordan	53, 93, 107
Babić, Tea	82	Horvat, Petar	96
Bakker, Eric	25	Ilić, Krunoslav	89
Ban, Željka	59	Inkret, Suzana	94
Bastić, Marijana	75	Ivanišević, Irena	95
Bauer, Jurica	76	Jakopin, Žiga	48
Bedeković, Nikola	66, 73, 82, 83, 90, 103	Janković, Patrizia	114
Bertone, Vlasta	104	Jasprica, Ivona	96, 119
Bertoša, Branimir	84	Javorić, Kristina	117
Bilandžić, Katijana	85	Jelić Matošević, Zoe	84
Biljan, Tomislav	37	Jeličić, Mario-Livio	97, 98, 110
Bjornstad, Vidar	96	Jozanović, Marija	57, 101, 112, 122, 123
Boček, Željka	99	Jurić-Babaj, Maja	74
Bohinc, Klemen	68	Kalafatović, Daniela	114
Bolanča, Tomislav	100	Kassal, Petar	95, 99, 121
Bonney, Karl	96	Kendall, Thomas	120
Bottema, Jeroen	42	Kirin, Srećko I.	65
Bregović, Nikola	80, 87, 96	Klačić, Tin	105
Budetić, Mateja	109	Klarić, Daniela Amidžić	97, 110
Bule, Kristina	100	Klarić, Ilija	97
Burić, Nataša	86	Knežević, Zdravka	60
Cappelletti, Giuseppe	49, 115	Kolympadi Markovic, Maria	112, 123
Čelan, Stipe	70	Košćak, Marta	59
Cinčić, Dominik	66, 83	Košutić Hulita, Nada	118
Cindro, Nikola	107, 110	Kovačević, Davor	68, 105
Cirino, Thomas	36	Kovačić, Jelena	97, 98, 110
Čiško-Anić, Blaženka	101	Kraikaew, Pitchnaree	25
Čonková, Miroslava	56, 108	Krivačić, Sara	99, 121
Čož-Rakovac, Rozelindra	123	Kučas, Filip	93
Crnolatac, Ivo	107	Kučić Grgić, Dajana	100
Cudic, Predrag	36	Kurpik, Gracjan	54
Cvetnić, Marija	87	Kurtanjek, Želimir	62
Cvetnić, Matija	100	Kuštro, Marija	100
Damala, Polyxeni	25	Lah, Jurij	28
Đambić, Kristina	122	Lazarenko, Viktoriia	80
Dandić, Andrea	109	Leko, Katarina	96
Danil de Namor, Angela F.	31	Letić, Nikolina	45
Della Ventura, Bartolomeo	57	Liović, Marin	66
Dodlek Šarkanj, Ivana	91	Logožar, Aleks	65
Domazet Jurašin, Darija	89	Lorincz, Artur	42
Dubaić, Karla	88	Lorincz, Kinga	42
Dutour Sikirić, Maja	69, 89, 94	Lovrić, Suzana	72
Edink, Ewald	42	Lubina, Katarina	122
Erceg, Ina	69, 89	Madunić-Čačić, Dubravka	57, 101, 122
Ernečić, Amalija	80	Magnabosco, Antonio	102
Forrest, Tara	25	Majchrzycki, Łukasz	56
Fotović, Luka	90	Majstorović, Gordana	102
Furač, Lidija	100	Majumder, Shahayra	36
Glumac, Nada	122	Makarić, Sandro	117
Golenić, Neven	93	Malčić, Mirta	72
Golenja, Josipa	91	Maltar Strmečki, Nadica	89, 94
Gomaz, Boris	92	Marder, Todd B.	59

Marić, Lucija	103
Marjanović, Nikola.....	118
Markić, Marinko	100
Markiewicz, Grzegorz	54, 79, 108
Marković, Dean.....	112, 122, 123
Maršavelski, Aleksandra	51
Martinis, Olgica.....	77
Martinović, Ivana.....	70
Maslov Bandić, Luna.....	39
Matijević, Gabrijela.....	123
Mavrinac, Marina	104
McLaughlin, Jay P.	36
Medančić, Tina	119
Mendeš, Davor	87
Mesić, Mia	105
Meštrović, Ernest.....	96, 119
Milašinović, Valentina	106
Milić, Marija.....	89
Miloloža, Martina	100
Mlinarić, Zvonimir	117
Modrušan, Matija.....	107
Molčanov, Krešimir.....	43, 106, 116
Mornar, Ana	97, 98
Mutavdžić Pavlović, Dragana.....	117
Nau, Werner M.....	24
Nussbaum, Robin.....	25
Odžak, Renata.....	85
Otmačić, Lucija	107
Pantalón Juraj, Natalija.....	65
Pargoletti, Eleonora.....	115
Patel, Swati A.....	120
Pavelić, Sandra Kraljević	123
Pavić, Monika	74
Perić, Berislav	65
Perlitiu, Filip	108
Peršić, Mateja.....	109
Piantanida, Ivo.....	59
Pičuljan, Katarina.....	82
Pilić, Zora	70
Portada, Tomislav.....	102
Požar, Josip	82, 90, 96
Prevarić, Viktorija	100
Pukleš, Iva	122
Rabuffetti, Marco	115
Radić, Vladimir.....	117
Radmilović, Nadežda	94
Rajić, Martina	110
Ranić, Jelena	119
Ravasio, Nicoletta.....	32
Rezić, Iva	40
Robescu, Marina Simona.....	115
Roca, Sunčica	80
Rolaniya, Asha	111
Safundžić Kučuk, Maša	60, 88
Sailani, Riya.....	63, 111
Sakač, Nikola	57, 101, 112, 122, 123
Samardžić, Mirela.....	109
Sangiorgio, Sara	115
Šarkanj, Bojan	91, 122
Saršon, Vesna.....	113
Šekutor, Marina	80, 81, 86
Semproli, Riccardo	115
Sepčić, Sabino	114
Sertić, Miranda.....	117
Silovski, Tajana	117
Siročić, Sandra.....	119
Skenderović, Anamarija	117
Smokrović, Kristina.....	94
Speranza, Giovanna	107, 115
Šprung, Matilda.....	85
Stacy, Heather.....	36
Stanić, Petra	116
Štefan, Leo	60, 88
Štefanić, Zoran	81, 92
Stefankiewicz, Artur R.	54, 56, 79, 108
Stejskal, Daria.....	71
Stilinović, Vladimir	46, 66, 83, 90, 103
Stolar, Tomislav.....	81
Štrbac, Petar.....	106
Sulyok, Michael	91
Šumonja, Josipa	113
Szántay, Csaba	27
Tavra, Dubravka	100
Tomišić, Vladislav	87, 96, 107
Tot, Dora	122
Turk, Ana Mornar	110
Turković, Lu	117
Ubiali, Daniela	115
Ukić, Šime.....	100
Usenik, Andrea.....	82
Užarević, Krunoslav.....	81
Velotta, Raffaele	57
Vianello, Robert	65, 96, 119
Vidović, Dorina	118
Vidović, Nikolina	107
Vinković Vrček, Ivana	89
Vranješević, Filip	112
Vrček, Valerije	112
Vuić, Juraj.....	117
Vukoja, Sanja.....	119
Walczak, Anna.....	54, 108
Williams, Brandon.....	36
Yadav, Jyoti	120
Zdrachek, Elena.....	25
Žgela, Diana.....	119
Živković, Igor	51
Zlatic, Gloria	70
Znc, Katarina	96
Zubak, Marko	95, 121
Zubčić, Snježana	97

PARTICIPANTS

Wiktorija Adamska
Marija Alešković
Jasna Alić Stolar
Melita Androlić
Klaudija Aušperger
Tea Babić
Eric Bakker
Mislav Barić
Marijana Bastić
Jurica Bauer
Nikola Bedeković
Ksenija Bedeković
Branimir Bertoša
Antonija Bešlić
Katijana Bilandžić
Rea Bilić
Tomislav Biljan
Ivana Blagović
Vesna Blažek Bregović
Lana Bogdanovski
Nikola Bregović
Neven Bublic
Mateja Budetić
Zrinka Buhin Šturlić
Giuseppe Cappelletti
Jurica Cebin
Nikola Cindro
Dražen Crnčec
Antonija Crnčec
Predrag Cudic
Marija Cvetnić
Vladimir Cvjetojević
Ana Čavlina
Kristina Činčurak
Miroslava Čonková
Nikolina Črnko Prpić
Vanja Čučković Makar
Biserka Ćurčija
Angela Danil de Namor
Aleksandar Danilovski
Dea Demeter
Petra Dičak
Ivana Došlović Pezić
Karla Dubaić
Matea Dujmić
Maximilian Dulson
Alen Đalapa
Ewald Edink
Ina Erceg
Borna Ferčec
Luka Fotović
Lucija Furjan
Andrea Gelemanović
Josipa Golenja
Boris Gomaz
Danijela Grabovac
Ivan Grgičević
Saša Grubešić
Carmen Guguta
Davor Hižak
Gordan Horvat
Petar Horvat
Suzana Inkret
Adriana Ivandić
Irena Ivanišević
Ivona Ivešić
Žiga Jakopin
Ivona Jasprica
Mario-Livio Jeličić
Ruža Jonjić
Maja Jurić-Babaja
Monika Kanižaj Silaj
Klara Karadakić
Dubravka Kardum
Petar Kassal
Thomas Kendall
Tetiana Khristova
Marin Knežević
Ankica Kolar
Lea Komočar
Jelena Kovač
Davor Kovačević
Sara Krivačić
Dubravko Kršul
Karla Kukina Gradečak
Gracjan Kurpik
Želimir Kurtanjek
Marija Kuštro
Marcel Lagedroste
Jurij Lah
Ivana Landeka
Viktoria Lazarenko
Katarina Leko
Nikolina Letić
Gabriela Lihtar
Marin Liović
Aleks Logožar
Marija Lovrić
Hrvojka Lukac
Adela Ljubić Jurišić
Dubravka Madunić-Čačić
Nina Magdalenić-Veselko
Antonio Magnabosco
Gordana Majstorović
Mirta Malčić
Lucija Marić
Nera Marjanović
Stanislav Martin

Valentina Martinček
Ivana Martinović
Luna Maslov Bandić
Maja Mataković
Marina Mavrinac
Tina Medančić
Karlo Meglič
Mia Mesić
Ernest Meštrović
Ines Mezga
Marijana Mihoci
Nina Mihoci
Maja Mikić Mašić
Mateja Miklečić Hendelja
Valentina Milašinović
Matija Modrušan
Krešimir Molčanov
Danijel Namjesnik
Werner Nau
Darija Navoj
Predrag Novak
Jasmina Novak
Jasminka Ostrički
Ivana Palej Jakopović
Monika Pavić
Marko Pavlović
Sanja Penić
Sara Perčić
Filip Perlitius
Mateja Peršić
Tomislav Petković
Emilija Petrović Hađar
Ivo Piantanida
Gordana Pintarić-Kovač
Tanja Poljak
Josip Požar
Dijana Premuš
Jasna Prnjavorac
Snježana Protulipac
Mirko Pupiće-Bakrač
Keith Racman
Elizabeta Rajič
Martina Rajić
Alma Ramić
Nicoletta Ravasio
Karla Remar
Iva Rezić
Rosana Ribić
Zrinka Runje Klaskan
Maša Safundžić Kučuk

Riya Sailani
Nikola Sakač
Mirela Samardžić
Vesna Saršon
Gaurav Saxena
Sabino Sepčić
Zlatko Smole
Giovanna Speranza
Petra Stanić
Snježana Starčević-Lovreković
Daria Stejskal
Vladimir Stilinović
Sanja Stipičević
Ingrid Stojko
Csaba Szántay, Jr.
Marina Šekutor
Martina Šeler Namjesnik
Olivera Šercer-Žokalj
Matija Šimac
Branko Špoljar
Anita Šporec
Vlado Štimac
Vitomir Šunjić
Zrinka Švob Troje
Dubravka Tavra
Patrik Tišma
Davor Tkalec
Vladislav Tomišić
Ines Topalović Piteša
Magda Topić
Mihaela Tuksar
Lu Turković
Andrea Usenik
Ivana Vadlja
Dorina Vidović
Nikolina Vidović
Adrijana Vinter
Mihaela Vrbnjak Grđan
Vesna Vrček
Sanja Vukoja
Elena Vukoša
Teo Vurnek
Dominic Welsh
Jyoti Yadav
Snježana Zdilar
Antun Zelić
Gloria Zlatic
Katarina Zrnc
Marko Zubak
Igor Živković



TERME SVETI MARTIN



VISIT sic.hkd.hr

**AND FOLLOW US ON SOCIAL MEDIA
FOR THE LATEST UPDATES & DETAILS**



  @SolutionsInChemistry #SiC2022
 @SiC_2022  Solutions in Chemistry